






## ORIGINAL RESEARCH ARTICLE

## Ecological Characterization of Malaria Vector Habitats in Awe and Nassarawa Eggon Local Government Areas, Nasarawa State, Nigeria

Victor Ameh Adejoh<sup>1</sup>, Akwashiki Ombugadu<sup>1</sup>, James Ishaku Maikenti<sup>1</sup>,  
Muhammed Ahmed Ashigar<sup>1</sup> and Victoria Adamu Pam<sup>1</sup>

<sup>1</sup>Department of Zoology, Faculty of Science, Federal University of Lafia, Nasarawa State, Nigeria

### ABSTRACT

Malaria remains a significant global health concern due to *Anopheles* mosquitoes. This study assessed malaria vector populations in Awe and Nassarawa Eggon LGAs of Nasarawa State, Nigeria. Mosquito larvae were collected employing standard dipping methods, while physicochemical parameters were analyzed in-situ and in the laboratory using standard methods. A total of 1,405 mosquito larvae were collected, with a higher abundance in Awe (95.2%) than in Nassarawa Eggon (4.8%). Culicines comprised 84.8% of the total larvae, while anophelines accounted for 15.2%. Although habitat type did not significantly influence *Anopheles* abundance ( $P = 0.364$ ), puddles (52.5%) and rice fields (28.6%) supported the highest larval populations. The highest mosquito abundance was recorded at a 351–400m gradient (47.8%). No significant association ( $P = 0.65$ ) was found between larval abundance and proximity to human dwellings. Temperature ( $r = 0.395$ ), electrical conductivity ( $r = 0.303$ ), salinity ( $r = 0.407$ ), total dissolved solids ( $r = 0.390$ ), and carbon dioxide ( $r = 0.015$ ) positively influenced *Anopheles* abundance, while pH ( $r = -0.039$ ), alkalinity ( $r = -0.024$ ), and dissolved oxygen ( $r = -0.067$ ) had negative correlations. *Anopheles gambiae* (87.2%) was more abundant than *An. funestus* (12.8%), with no significant difference ( $P = 0.0881$ ) between species in both LGAs. Continued mosquito surveillance in Awe and Nassarawa Eggon is necessary to enhance vector control strategies and malaria prevention efforts.

### ARTICLE HISTORY

Received December 27, 2024

Accepted March 27, 2025

Published March 31, 2025

### KEYWORDS

*Anopheles*, Habitat, Malaria, Mosquito, Nasarawa



© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

### INTRODUCTION

*Anopheles* species have plagued the world with malaria for centuries. In 2021, approximately 247 million malaria cases were reported across 84 malaria-endemic countries, marking an increase from 245 million cases in 2020 (WHO, 2022). Twenty-nine countries accounted for 96% of malaria cases globally, and four countries – Nigeria (30.9%), the Democratic Republic of Congo (11.3%), Niger (5.9%), and United Republic of Tanzania (4.3%) – accounted for almost half of all cases globally (WHO, 2022, 2024).

The distribution and population size of adult *Anopheles* mosquitoes depend on the availability and productivity of their larval breeding sites (Lapang *et al.*, 2019; Oniya *et al.*, 2019). The productivity of such habitats influences the conditions of larval development and plays a major role in the eventual size of adult mosquitoes, which in turn affects factors such as longevity, abundance, morphology, and blood meal volume (Ukubuiwe *et al.*, 2018; Hassan *et al.*, 2021; Njila *et al.*, 2023). While previous studies, such as Lapang *et al.* (2019) and Hassan *et al.* (2021), have

explored habitat productivity and vector abundance, their findings were limited to specific ecological zones, leaving gaps in understanding how varying environmental conditions influence vector populations across different landscapes. Similarly, Oguiche *et al.* (2022) assessed malaria vector ecology but did not comprehensively incorporate spatial mapping techniques to visualize habitat distribution and vector density.

Building spatial models of disease incidence and risk, such as maps for the possible spread of infectious diseases like malaria, is a key theoretical objective of epidemiology. In particular, Ryan *et al.* (2020) emphasize risk maps for malaria in Africa. Unfortunately, there is little information on the availability of risk maps for malaria vectors, particularly *Anopheles* species, in or around Awe and Nassarawa Eggon LGAs, Nasarawa State, Nigeria. This study addresses these gaps by integrating ecological characterization with geospatial mapping to provide a more comprehensive understanding of malaria vector habitats in the study areas.

**Correspondence:** Victor Ameh Adejoh. Department of Zoology, Faculty of Science, Federal University of Lafia, Nasarawa State, Nigeria. PMB 146 Makurdi Road, Gandu, Lafia Nasarawa State, Nigeria. ✉ [amehvictor02@gmail.com](mailto:amehvictor02@gmail.com).

**How to cite:** Adejoh, V. A., Ombugadu, A., Maikenti, J. I., Ashigar, M. A., & Pam, V. A. (2025). Ecological Characterization of Malaria Vector Habitats in Awe and Nassarawa Eggon Local Government Areas, Nasarawa State, Nigeria. *UMYU Scientifica*, 4(1), 317 – 324. <https://doi.org/10.56919/usci.2541.031>

**MATERIALS AND METHODS**

**Study Area**

The study was carried out in Awe and Nassarawa Eggon LGAs, respectively, in Nasarawa State, Nigeria (Figure 1).

**Identification and Characterization of Anopheles Breeding Habitat**

A preliminary survey was carried out prior to the fieldwork to identify potential *Anopheles* mosquito breeding habitats. The respective coordinates of identified breeding habitats were obtained with the aid of a Garmin GPS navigator (Garmin Inc., Kansas, USA) and exported to ArcGIS version 9.3, which was then used to generate a map of the study areas (Egwu *et al.*, 2018; Njila *et al.*, 2023).


Sampling was conducted between 7:00 am and 11:00 am (Egwu *et al.*, 2018; Lapang *et al.*, 2019). All potential mosquito breeding sites were carefully inspected for the presence of *Anopheles* larvae through direct visual observation (Mattah *et al.*, 2017; Lapang *et al.*, 2019). Habitats of the *Anopheles* mosquito species were categorized with respect to whether they are man-made or natural (Owolabi & Bagbe, 2020).

**Sample Collection**

*Anopheles* mosquito larvae were sampled using the standard dipping technique as described by Oniya *et al.* (2019). At each identified breeding site, a 350 mL standard dipper was used to collect larvae by submerging it at a 45-degree angle into the water. In shallow or vegetated habitats, the dipper was skimmed gently along the surface to avoid disturbing the larvae. A total of 10 dips were taken per site. Collected larvae were transferred into labelled plastic containers filled with site water and covered with fine mesh netting for transport to the laboratory.

**Determination of Physicochemical Parameters**

Temperature, pH, electrical conductivity (EC), salinity, and total dissolved solids (TDS) were measured in situ using a well-calibrated Oakton Multi-Parameter (Oakton, USA), following the protocol described by Lin *et al.* (2021). Measurements were taken directly at each sampling site by immersing the probe into the water, allowing for real-time readings.



**Figure 1: Map of Awe and Nassarawa Eggon LGAs Showing Mosquito Larvae Sample Points**

Alkalinity (mg/L CaCO<sub>3</sub>), carbon dioxide (CO<sub>2</sub>), and dissolved oxygen (DO<sub>2</sub>) were determined in the

laboratory following standard titrimetric and electrochemical methods as described by Pimenta *et al.*

(2023). Water samples were collected in airtight polyethylene bottles to minimize gas exchange and contamination. Samples were transported in cool, insulated containers to maintain stability and prevent alterations in chemical composition. Alkalinity was measured using acid-base titration, CO<sub>2</sub> levels were determined by acidification followed by titration, and DO<sub>2</sub> was quantified using the Winkler method (Hu *et al.*, 2024). All analyses were conducted immediately upon arrival at the laboratory to ensure accuracy and reliability of results.

**Anopheles Mosquito Rearing**

The larvae were identified based on their morphological characteristics (Gillies & Coetzee, 1987; Sallum *et al.*, 2020). All larvae morphologically confirmed as belonging to *Anopheles* species were used for this study. The larvae were reared under controlled conditions following standard procedures (Zengenene *et al.*, 2022). They were housed in plastic rearing cages (30 cm × 25 cm × 5 cm) and maintained in larval bowls filled with water sourced from their respective natural habitats to preserve their native environmental conditions. This setup ensured optimal development by minimizing stress and maintaining natural microbial communities essential for larval growth. Water levels were monitored and replenished as needed to prevent desiccation, while environmental factors such as temperature and light exposure were kept stable to support normal larval development (Wohl & McMeniman, 2022).

Emerged adult *Anopheles* mosquitoes were fed with 10% sucrose solution (Ukubuiwe *et al.*, 2018; Lapang *et al.*, 2019; Ngowo *et al.*, 2021), knocked down in a -20°C freezer for 10 minutes, and then further identified under a dissecting

microscope using identification keys by Ukubuiwe *et al.*, (2018) and Coetzee (2020).

**Data Analysis**

The data was analysed using R statistical software (version 3.6.1). One-Way ANOVA was applied to compare larval abundance across different breeding habitats and distances from the nearest houses. A Chi-Square test was used to assess differences in *Anopheles* mosquito larvae between the two LGAs. Descriptive statistics provided mean values and percentages of larval abundance in the study sites. Pearson’s Correlation Coefficient measured the relationship between larval abundance and physicochemical parameters, while another Chi-Square test examined the association between emerged adult *Anopheles* species in both LGAs. A P-value below 0.05 was considered statistically significant.

**RESULTS AND DISCUSSION**

**Association between Physicochemical Parameters and Anopheline Larval Abundance in Awe and Nassarawa Eggon, Nigeria**

There was a positive relationship between temperature ( $r = 0.395$ ), electrical conductivity ( $r = 0.303$ ), salinity ( $r = 0.407$ ), total dissolved solid ( $r = 0.390$ ), and carbon dioxide ( $r = 0.015$ ) and the abundance of anopheline mosquito larvae in their breeding habitats at Awe and Nassarawa Eggon LGA, Nasarawa State, Nigeria. Conversely, pH ( $r = -0.039$ ), alkalinity ( $r = -0.024$ ), and dissolved oxygen ( $r = -0.067$ ) all recorded a negative relationship with the abundance of anopheline mosquito larvae in their breeding habitats at Awe and Nassarawa Eggon LGA, Nasarawa State, Nigeria (Table 1).

**Table 1: Association between Physicochemical Parameters and Anopheline Larval Abundance in Awe and Nassarawa Eggon, Nigeria**

| Physicochemical Parameters | P     | r      | Coefficient of Determination (r <sup>2</sup> ) |
|----------------------------|-------|--------|--|
| Temperature                | 0.094 | 0.395  | 0.156025                                       |
| pH                         | 0.874 | -0.039 | 0.001521                                       |
| Electrical Conductivity    | 0.207 | 0.303  | 0.091809                                       |
| Salinity                   | 0.084 | 0.407  | 0.165649                                       |
| Total Dissolved Solids     | 0.098 | 0.390  | 0.15210  |
| Alkalinity                 | 0.921 | -0.024 | 0.000576                                       |
| Carbon Dioxide             | 0.952 | 0.015  | 0.000225                                       |
| Dissolved Oxygen           | 0.785 | -0.067 | 0.004489                                       |

The negative correlation observed in the relationship between anopheline mosquito larvae abundance and pH, alkalinity, and dissolved oxygen possibly suggests that they do not influence the abundance of anopheline larvae in habitats sampled in Awe and Nassarawa Eggon LGA, Nasarawa State, Nigeria. This result is consistent with that of Obi *et al.* (2019), Akeju *et al.* (2022), and Seal and Chatterjee (2023), who all recorded a negative correlation in the relationship between anopheline mosquito larvae abundance and pH, alkalinity, and dissolved oxygen in their respective study sites.

The positive correlation observed between Electrical conductivity (EC), salinity, total dissolved solid (TDS),

and carbon dioxide (CO<sub>2</sub>) contents of the water from larval habitats and anopheline larval abundance at the study locations suggests that these parameters influence the abundance of these anophelines. Oguche *et al.* (2022) and Omondi *et al.* (2023) reported similar associations with EC, salinity, TDS, CO<sub>2</sub> and DO<sub>2</sub> content with anophelines and culicines in their study in Kaduna State, Nigeria, and Kwale County, Kenya, respectively.

**Checklist of Field Collected Larvae in Awe and Nassarawa Eggon LGAs of Nasarawa State, Nigeria**

A total of 1,405 mosquito larvae belonging to the anopheline and culicine groups were collected from various habitats in Awe (95.2%) and Nassarawa Eggon

(4.8%) LGAs, respectively. Culicine larvae (84.7%) were the predominant group encountered, while the anophelines accounted for 15.2% of the total larval population collected from the field (Table 2). The abundance of mosquito larvae species differed significantly ( $\chi^2 = 265.81$ ,  $df = 1$ ,  $P < 0.0001$ ) between Awe and Nasarawa Eggon LGAs, Nasarawa State, Nigeria, indicating a notable variation in species distribution across the study locations. This study revealed high larval heterogeneity, suggesting that environmental factors such as habitat type and water quality influence mosquito breeding site selection. The observed differences may also be attributed to ecological variations between the two LGAs, such as differences in vegetation cover, human activities, and climatic conditions, which create distinct microhabitats that support varying mosquito species compositions. These

results are consistent with studies by Lapang *et al.* (2019) in Shendam, Ekedo *et al.* (2020) in Umudike, Nigeria, and Oforika *et al.* (2024) in Lagos State, Nigeria. Conversely, Surendran *et al.* (2021) reported mosquito larvae productivity as a habitat species specific in Jaffna Peninsula, Sri Lanka.

The great variation observed in larval abundance between anopheline and culicine mosquitoes could indicate that the habitat types studied possibly support more of the growth of culicines than the anophelines. This result is in tandem with that of Alkhayat *et al.* (2020) in Qatar, Gowelo *et al.* (2020) in Malawi, and Onen *et al.* (2021) in Central Uganda. However, the result did not agree with that of Lapang *et al.* (2019) and Kinga *et al.* (2022), who recorded no variation in larval abundance between anophelines and culicines in their study.

**Table 2: Checklist of Field Collected Larvae in Awe and Nassarawa Eggon LGAs of Nasarawa State, Nigeria**

| Group      | Awe LGA (%) |             | Nassarawa Eggon LGA (%) |            | Total       |      |
|------------|-------------|-------------|-------------------------|------------|-------------|------|
|            | Number      | (%)         | Number                  | (%)        | Number      | (%)  |
| Anopheline | 157         | 11.7        | 57                      | 85.1       | 214         | 15.2 |
| Culicine   | 1181        | 88.3        | 10                      | 14.9       | 1191        | 84.8 |
|            | <b>1338</b> | <b>95.2</b> | <b>67</b>               | <b>4.8</b> | <b>1405</b> |      |

***Emerged Adult Anopheles Mosquitoes***

Only 164 of the 214 *Anopheles* mosquito species collected from their natural habitats emerged. Of this number, *Anopheles gambiae* (87.2%) was most abundant, while *An. funestus* accounted for 12.8% of the population (Table 3). The distribution of emerged *Anopheles* mosquito species between Awe and Nasarawa Eggon LGAs, Nasarawa State, Nigeria, showed no statistically significant variation ( $\chi^2 = 2.9086$ ,  $df = 1$ ,  $P = 0.0881$ ). This indicates that the observed differences in species composition between the two LGAs could be due to random variation rather than distinct ecological or environmental factors. The relatively high *P*-value suggests that *Anopheles* species are likely

distributed uniformly across both locations, potentially influenced by similar habitat characteristics, breeding site availability, and climatic conditions. The consistent presence of *Anopheles* mosquitoes in this study suggests an ongoing risk of malaria transmission in the area. The dominance of *An. gambiae* may be due to its remarkable adaptability, allowing it to thrive even in challenging environments. This aligns with findings by Hassan *et al.* (2021) and Li *et al.* (2021), who also observed its resilience in adverse conditions. This result is consistent with that of Lapang *et al.* (2019) and Osidoma *et al.* (2023), where they had more emergence of *An. gambiae* than other mosquito species reared.

**Table 3: Emerged Adult Anopheles Mosquitoes**

| LGA              | No. of Emerged Adult Mosquitoes |                  |                  |                    |                 |                 | Overall Emerged (%) |
|------------------|---------------------------------|------------------|------------------|--------------------|-----------------|-----------------|---------------------|
|                  | <i>A. gambiae s. l.</i>         |                  | Total (%)        | <i>A. funestus</i> |                 | Total (%)       |                     |
|                  | Male                            | Female           |                  | Male               | Female          |                 |                     |
| Awe              | 27(25.5)                        | 79(74.5)         | <b>106(74.1)</b> | 5(45.5)            | 6(54.5)         | <b>11(52.4)</b> | <b>117(71.3)</b>    |
| Nassarawa Eggon  | 7(18.9)                         | 30(81.1)         | <b>37(25.9)</b>  | 3(27.3)            | 7(70.0)         | <b>10(47.6)</b> | <b>47(28.7)</b>     |
| <b>Total (%)</b> | <b>34(23.8)</b>                 | <b>109(76.2)</b> | <b>143(87.2)</b> | <b>8(38.1)</b>     | <b>13(61.9)</b> | <b>21(12.8)</b> | <b>164</b>          |

***Distribution of Anopheline Mosquito Larvae Across Different Habitat Types in Awe and Nasarawa Eggon LGAs, Nasarawa State***

Anopheline Mosquito larvae were most abundant in puddle habitat (52.5%) habitat followed by rice field (28.6%), while the animal hoofs (0.2%) accounted for the least population (Figure 2). However, there was no significant difference ( $F = 1.37$ ,  $df = 4$ ,  $Adj.R^2 = 13.98\%$ ,  $P = 0.364$ ) in the abundance of anopheline mosquito larvae in Awe and Nassarawa Eggon LGA, Nasarawa State, Nigeria. The absence of significant variations in the abundance of anopheline mosquito larvae across different habitat types in Awe and Nassarawa Eggon LGA of

Nasarawa State, Nigeria, strongly indicates that both locations provide favorable conditions for breeding this disease-carrying vector. The abundance of mosquito larvae in puddle habitats can be attributed to the rainy season, which creates numerous pockets of puddle habitats favorable for mosquitoes.

This finding aligns with the research conducted by Ekedo *et al.* (2020), and Getachew *et al.* (2020). It did not agree with the results of Lapang *et al.* (2019), who collected more mosquito larvae in rice fields at Shendam, and Oguche *et al.* (2022), who collected more mosquito larvae from abandoned tyres in Kaduna metropolis.

**Distribution of Anopheline Mosquito Larvae Based on Proximity of Breeding Sites to Residential Areas in Awe and Nasarawa Eggon LGAs, Nasarawa State, Nigeria**

The gradient 351–400 m recorded the highest population of *Anopheles* mosquito larvae (47.8%), followed by the 251–300 m gradient (14.3%), while the 451–500 m gradient had the least abundance (0.4%) (Figure 3). This pattern suggests that *Anopheles* mosquitoes preferentially breed within specific elevation ranges, possibly due to favorable environmental conditions such as water availability, vegetation cover, and reduced disturbance. However, statistical analysis showed no significant difference in larval abundance relative to the proximity of breeding habitats to human dwellings ( $F = 0.76$ ,  $df = 9$ ,  $Adj. R^2 = 0.000\%$ ,  $P = 0.65$ ). This indicates that the

distance between breeding sites and human settlements does not directly influence *Anopheles* larval density. The high abundance observed at 351–400 m suggests that *Anopheles* mosquitoes are well adapted to breeding even at moderate distances from human habitations, reinforcing the need for vector control strategies that extend beyond immediate residential areas.

These findings align with the observations made by Lapang *et al.* (2019) and Zogo *et al.* (2019), who reported greater larval abundance in habitats surrounded by fewer houses and more green areas compared to habitats densely populated by human inhabitants. However, the result is in contrast with that of Mathania *et al.* (2020), who recorded more abundance of anopheline larvae in habitats along gradients ranging between 51-100m in Tanzania.




Figure 2: Distribution of Anopheline Mosquito Larvae Across Different Habitat Types in Awe and Nasarawa Eggon LGAs, Nasarawa State




Figure 3: Distribution of Anopheline Mosquito Larvae Based on Proximity of Breeding Sites to Residential Areas in Awe and Nasarawa Eggon LGAs, Nasarawa State, Nigeria

**Breeding Index (BI) of Anopheline Larvae Across Different Habitat Types in Awe and Nasarawa Eggon LGAs, Nasarawa State, Nigeria**

The puddle habitat proved to be the most favorable breeding ground, recording a breeding index (BI) of 31.7, followed by the rice field (6.6), while the ditches (0.7) had the least BI (Table 4). The animal hoof prints had a BI of 0.0, indicating that they did not contribute to mosquito breeding in the study area. The consistently high BI observed in puddles suggests that these habitats provide ideal conditions for *Anopheles* larvae, possibly due to factors such as stagnant water, suitable temperature, and organic matter availability, which enhance larval survival and development. With an overall breeding index of 8.2,

which exceeds the 5.0 threshold for high mosquito productivity, the *Anopheles* population in Awe and Nasarawa Eggon LGAs presents a serious public health risk. The elevated BI suggests an increased likelihood of multiple mosquito-borne disease outbreaks among residents, necessitating urgent vector control interventions. These findings align with the studies of Doumbe-Belisse *et al.* (2021) and Chan *et al.* (2022), who reported even higher breeding indices of 20.0 and 48.8, respectively, reinforcing the idea that high BI values correspond to greater malaria transmission potential. The observed differences in BI across habitats further emphasize the need for targeted control measures to mitigate malaria transmission in the study area, focusing on high-risk breeding sites such as puddles.

**Table 4: Breeding Index (BI) of Anopheline Larvae Across Different Habitat Types in Awe and Nasarawa Eggon LGAs, Nasarawa State, Nigeria**

| Habitats          | Communities |          |            |            |            |            |                    |            |            | Total BI    |
|-------------------|-------------|----------|------------|------------|------------|------------|--------------------|------------|------------|-------------|
|                   | Awe LGA     |          |            |            |            |            | Nasarawa Eggon LGA |            |            |             |
|                   | KA          | MG       | MSM        | KKR        | TG         | Sub-Total  | NEQM               | NEM        | Sub-Total  |             |
| Puddle            | 0.4         | 0.0      | 8.7        | 9.6        | 9.3        | <b>28</b>  | 1                  | 2.7        | <b>3.7</b> | <b>31.7</b> |
| Rice Field        | 0.0         | 4.8      | 0.0        | 0.0        | 0.0        | <b>4.8</b> | 1.8                | 0.0        | <b>1.8</b> | <b>6.6</b>  |
| Ditch             | 0.0         | 0.0      | 0.0        | 0.7        | 0.0        | <b>0.7</b> | 0.0                | 0.0        | <b>0.0</b> | <b>0.7</b>  |
| Animal Hoof       | 0.0         | 0.0      | 0.0        | 0.0        | 0.0        | <b>0.0</b> | 0.0                | 0.0        | <b>0.0</b> | <b>0.0</b>  |
| Tyre Track        | 0.0         | 0.0      | 0.0        | 0.0        | 0.0        | <b>0.0</b> | 0.5                | 1.7        | <b>2.2</b> | <b>2.2</b>  |
| <b>Average BI</b> | <b>0.1</b>  | <b>1</b> | <b>1.7</b> | <b>2.1</b> | <b>1.9</b> | <b>6.7</b> | <b>0.7</b>         | <b>0.9</b> | <b>1.5</b> | <b>8.2</b>  |

KA: Kauje-Aburi, MG: Manga-Galadima, MSM: Madiki Salt Mine, KKR: Kekura, TG: Tunga, NEQM: Nasarawa Eggon Quarry Mine, NEM: Nasarawa Eggon Market

**CONCLUSION**

This study clearly shows the presence of heterogenous association between mosquito larval groups (anopheline and culicine) across breeding habitats in Awe and Nassarawa Eggon LGAs, Nasarawa State, Nigeria. Emerged adult *Anopheles gambiae* complex had a higher abundance than *A. funestus*. All the mosquito larvae collected utilized five habitat types: animal hoof, ditch, puddle, rice fields, and tyre tracks.

Also, this research shows that mosquitoes have no preferred gradient from houses. However, on average, they preferred to breed farther away from human settlements. Furthermore, it was observed that physicochemical parameters (temperature, EC, salinity, TDS & CO<sub>2</sub>) positively influence larval abundance in the study areas with the exception of pH, alkalinity, and DO<sub>2</sub>. The observed high breeding index (8.2) recorded in this study suggests that the inhabitants of Awe and Nassarawa Eggon LGA are at risk of the transmission of mosquito-borne diseases.

Mosquito surveys should be carried out all year round in Awe and Nassarawa Eggon LGAs. This will provide additional information on their distribution within these areas. Inhabitants in these areas should always clear stagnant water bodies so as to interrupt mosquito breeding success. Residents of the two LGAs should sleep under insecticide-treated bed nets in order to avoid human-vector contact.

**ACKNOWLEDGEMENT**

We sincerely appreciate the residents of Awe and Nasarawa Eggon LGAs for their kindness and cooperation in allowing us to conduct this field survey.

**REFERENCES**

Akeju, A. V., Olusi, T. A., & Simon-Oke, I. A. (2022). Effect of physicochemical parameters on *Anopheles* mosquitoes larval composition in Akure North Local Government Area of Ondo State, Nigeria. *The Journal of Basic and Applied Zoology*, 83(1), 34. [\[Crossref\]](#)

Alkhatay, F. A., Ahmad, A. H., Rahim, J., Dieng, H., Ismail, B. A., Imran, M., Sheikh, U. A., Shahzad, M. S., Abid, A. D., & Munawar, K. (2020). Characterization of mosquito larval habitats in Qatar. *Saudi Journal of Biological Sciences*, 27(9), 2358-2365. [\[Crossref\]](#)

Chan, K., Tusting, L. S., Bottomley, C., Saito, K., Djouaka, R., & Lines, J. (2022). Malaria transmission and prevalence in rice-growing versus non-rice-growing villages in Africa: A systematic review and meta-analysis. *The Lancet Planetary Health*, 6(3), e257-e269. [\[Crossref\]](#)

Coetzee, M. (2020). Key to the females of Afrotropical *Anopheles* mosquitoes (*Diptera*: Culicidae). *Malaria Journal*, 19(1), 70. [\[Crossref\]](#)

Dikin, S. K., Schuster-Wallace, C. J., & Elliot, S. J. (2011). Developing a vulnerability mapping methodology: Applying the water-associated disease index to

- dengue in Malaysia. *PLoS ONE*, 8(5), Article e63584. [\[Crossref\]](#)
- Dondorp, A. M., Nosten, F., Yi, P., Das, D., Physo, A. P., Tarning, J., Lwin, K. M., Arley, F., Hanpithakpong, W., Lee, S. J., Ringwald, P., Silamut, K., Imwong, M., Chotivanich, K., Lim, P., Herdman, T., An, S. S., Yeung, S., Singhasivanon, P., ... White, N. J. (2009). Artemisinin resistance in *Plasmodium falciparum* malaria. *New England Journal of Medicine*, 361(5), 455-467. [\[Crossref\]](#)
- Doumbe-Belisse, P., Kopya, E., Ngadjjeu, C. S., Sonhafouo-Chiana, N., Talipouo, A., Djamouko-Djonkam, L., Awono-Ambene, P. H., Wondji, C. S., Njiokou, F., & Antonio-Nkondjio, C. (2021). Urban malaria in sub-Saharan Africa: Dynamic of the vectorial system and the entomological inoculation rate. *Malaria Journal*, 20(1), 364. [\[Crossref\]](#)
- Egwu, O. A., Ohaeri, C. C., Amaechi, E. C., & Ehisianya, C. N. (2018). Distribution and abundance of mosquito larvae in Ohafia, Abia State, Nigeria. *Cuadernos de Investigación UNED*, 10(2), 379-385. [\[Crossref\]](#)
- Ekedo, C. M., Okore, O. O., Uzoma, V. C., & Okoro, P. I. (2020). Habitat preference of mosquito larvae in Michael Okpara University of Agriculture, Umudike, Nigeria. *International Journal of Mosquito Research*, 7(4), 6-9.
- Ferede, G., Tiruneh, M., Abate, E., Kassa, W. J., Wondimeneh, Y., Damtie, D., & Tessema, B. (2018). Distribution and larval breeding habitats of *Aedes* mosquito species in residential areas of northwest Ethiopia. *Epidemiology and Health*, 40, Article e2018015. [\[Crossref\]](#)
- Getachew, D., Balkew, M., & Tekie, H. (2020). *Anopheles* larval species composition and characterization of breeding habitats in two localities in the Ghibe River Basin, southwestern Ethiopia. *Malaria Journal*, 19(1), 65. [\[Crossref\]](#)
- Gillies, M. T., & Coetzee, M. (1987). *A supplement to the Anophelinae of Africa south of the Sahara (Afrotropical region)*. South African Institute for Medical Research.
- Gowelo, S., Chirombo, J., Koenraadt, C. J. M., Mzilahowa, T., van den Berg, H., Takken, W., & McCann, R. S. (2020). Characterisation of anopheline larval habitats in southern Malawi. *Acta Tropica*, 212, Article 105558. [\[Crossref\]](#)
- Hassan, S. C., Olayemi, I. K., Omalu, I. C. J., Adefolalu, F. S., Eke, S. S., & Otuu, C. A. (2021). Spatiotemporal distribution and composition of *Anopheles* mosquito species in some selected eco-settings of Nasarawa State North Central Nigeria. *American Journal of Biology and Life Sciences*, 9(1), 1-9.
- Hu, Y., Xu, Y., Wang, H., Ren, H., & He, Y. (2024). Assessment of an automated titration system for batch measurements of total alkalinity. *Marine Chemistry*, 260, Article 104435. [\[Crossref\]](#)
- Kinga, H., Kengne-Ouafo, J. A., King, S. A., Egyirifa, R. K., Aboagye-Antwi, F., & Akorli, J. (2022). Water physicochemical parameters and microbial composition distinguish *Anopheles* and *Culex* mosquito breeding sites: Potential as ecological markers for larval source surveillance. *Journal of Medical Entomology*, 59(5), 1817-1826. [\[Crossref\]](#)
- Lapang, P. M., Ombugadu, A., Ishaya, M., Mafuyai, M. J., Njila, H. L., Nkup, C. D., & Mwansat, G. S. (2019). Abundance and diversity of mosquito species larvae in Shendam LGA, Plateau State, North-Central Nigeria: A panacea for vector control strategy. *Journal of Zoological Research*, 3(3), 25-33. [\[Crossref\]](#)
- Li, C., Gao, Y., Zhao, Z., Dang, D., Zhou, R., Wang, J., Zhang, Q., & Liu, Q. (2021). Potential geographical distribution of *Anopheles gambiae* worldwide under climate change. *Journal of Biosafety and Biosecurity*, 3(2), 82-89. [\[Crossref\]](#)
- Lin, J., Tsai, H., & Lyu, W. (2021). An integrated wireless multi-sensor system for monitoring the water quality of aquaculture. *Sensors*, 21(24), 8179. [\[Crossref\]](#)
- Mahgoub, M. M., Kweka, E. J., & Himeidan, Y. E. (2017). Characterization of larval habitats, species composition and factors associated with the seasonal abundance of mosquito fauna in Gezira, Sudan. *Infectious Diseases of Poverty*, 6(1), 23. [\[Crossref\]](#)
- Mathania, M. M., Munisi, D. Z., & Silayo, R. S. (2020). Spatial and temporal distribution of *Anopheles* mosquito's larvae and its determinants in two urban sites in Tanzania with different malaria transmission levels. *Parasite Epidemiology and Control*, 11, Article e00179. [\[Crossref\]](#)
- Mattah, P. A. D., Futagbi, G., Amekudzi, L. K., Mattah, M. M., de Souza, D. K., Kartie-Attipoe, W. D., Bimi, L., & Wilson, M. D. (2017). Diversity in breeding sites and distribution of *Anopheles* mosquitoes in selected urban areas of southern Ghana. *Parasites & Vectors*, 10(1), 25. [\[Crossref\]](#)
- Ngowo, H. S., Hape, E. E., Matthiopoulos, J., Ferguson, H. M., & Okumu, F. O. (2021). Fitness characteristics of the malaria vector *Anopheles funestus* during an attempted laboratory colonization. *Malaria Journal*, 20(1), 148. [\[Crossref\]](#)
- Njila, H. L., Sani, H. S., Ombugadu, A., & Tanko, N. S. (2023). Evaluation of coloured containers as attractants to gravid mosquitoes for oviposition. *Trends in Technical & Scientific Research*, 6(1), 1-8. [\[Crossref\]](#)
- Obi, O. A., Nock, I. H., & Adebote, D. A. (2019). Biodiversity of microinvertebrates co-inhabiting mosquitoes habitats in patchy rock pools on inselbergs within Kaduna State, Nigeria. *The Journal of Basic and Applied Zoology*, 80(1), 57. [\[Crossref\]](#)
- Oforika, C., Omotayo, A., & Adeleke, M. (2024). Seasonal diversity in mosquito larval ecology and its public health implications in urban slums of Lagos, Nigeria. *The American Journal of Tropical Medicine and Hygiene*. Advance online publication. [\[Crossref\]](#)
- Oguiche, O. D., Auta, I. K., Ibrahim, B., Yayock, H. C., & Johnson, O. (2022). Breeding sites characteristics and mosquito abundance in some selected locations

- within Kaduna metropolis. *FUDMA Journal of Sciences*, 6(6), 70-75. [\[Crossref\]](#)
- Okoh, H. I., Akinfemiwa, C. A., Iyeh, C., Mogaji, H. O., Omonijo, A. O., & Awe, O. B. (2020). Spatial distribution and breeding habitat characterization of *Anopheles* mosquitoes in Ado-Ekiti, Ekiti State, Nigeria. *FUOYE Journal of Pure and Applied Sciences*, 5(1), 154-164.
- Okorie, P. N., McKenzie, F. E., Ademowo, O. G., Bockarie, M., & Kelly-Hope, L. (2011). Nigeria *Anopheles* vector database: An overview of 100 years' research. *PLoS ONE*, 6(12), Article e28347. [\[Crossref\]](#)
- Omondi, C. J., Nonoh, J., & Ntabo, R. (2023). Characterization of *Anopheles funestus* larval habitats in Fiyoni, Kwale County, Kenya: Insights into malaria vector ecology and control. *European Journal of Ecology*, 9(2). [\[Crossref\]](#)
- Onen, H., Odong, R., Chemurot, M., Tripet, F., & Kayondo, J. K. (2021). Predatory and competitive interaction in *Anopheles gambiae* sensu lato larval breeding habitats in selected villages of central Uganda. *Parasites & Vectors*, 14(1), 331. [\[Crossref\]](#)
- Oniya, M. O., Adeyekun, A. L., & Olusi, T. A. (2019). Ecological factors favoring mosquito breeding in Ifedore Local Government Area of Ondo State, Nigeria. *Journal of Ecology and the Natural Environment*, 11(6), 68-74. [\[Crossref\]](#)
- Osidoma, E., Pam, V., Uzoigwe, N., Ombugadu, A., Omalu, I., Maikenti, J., Attah, A., Ashigar, M., & Dogo, S. (2023). A study on mosquitoes composition and malaria transmission in some communities in Doma Local Government Area of Nasarawa State, Nigeria. *Journal of Bioscience and Biotechnology Discovery*, 8(1). [\[Crossref\]](#)
- Owolabi, D. O., & Bagbe, A. S. (2020). Assessment of physico-chemical and ecological variables in selected natural breeding sites of mosquitoes in Ibadan, Oyo State, Nigeria. *Archives of Pharmacy & Pharmacology Research*, 1(5), APPR.MS.ID.000521. [\[Crossref\]](#)
- Pimenta, A. R., Oczkowski, A. J., McKinney, R. A., & Grear, J. S. (2023). Geographical and seasonal patterns in the carbonate chemistry of Narragansett Bay, RI. *Regional Studies in Marine Science*, 62, Article 102903. [\[Crossref\]](#)
- Rice, E. W., Baird, R. B., Eaton, A. D., & Clesceri, L. S. (Eds.). (2012). *Standard methods for the examination of water and wastewater* (22nd ed.). American Public Health Association.
- Ryan, S. J., Lippi, C. A., & Zermoglio, F. (2020). Shifting transmission risk for malaria in Africa with climate change: A framework for planning and intervention. *Malaria Journal*, 19(1), 170. [\[Crossref\]](#)
- Sallum, M. A. M., Obando, R. G., Carrejo, N., & Wilkerson, R. C. (2020). Identification keys to the *Anopheles* mosquitoes of South America (Diptera: Culicidae). I. Introduction. *Parasites & Vectors*, 13(1), 583. [\[Crossref\]](#)
- Seal, M., & Chatterjee, S. (2023). Combined effect of physico-chemical and microbial quality of breeding habitat water on oviposition of malarial vector *Anopheles subpictus*. *PLoS ONE*, 18(3), Article e0282825. [\[Crossref\]](#)
- Snow, R. W., Marsh, K., & Le Sueur, D. (1996). The need for maps of transmission intensity to guide malaria control in Africa. *Parasitology Today*, 12(11), 455-457. [\[Crossref\]](#)
- Surendran, S. N., Jayadas, T. T. P., Thiruchenthooran, V., Raveendran, S., Tharsan, A., Santhirasegaram, S., Sivabalakrishnan, K., Karunakaran, K., Ponnaiah, B., Gomes, L., Malavige, G. N., & Ramasamy, R. (2021). *Aedes* larval bionomics and implications for dengue control in the paradigmatic Jaffna peninsula, northern Sri Lanka. *Parasites & Vectors*, 14(1), 162. [\[Crossref\]](#)
- Ukubuiwe, A. C., Olayemi, I. K., Arimoro, F. O., Omalu, I. C. J., Baba, B. M., Ukubuiwe, C. C., Odeyemi, M. O., & Adeyemi, K. A. (2018). Influence of rearing-water temperature on life stages' vector attributes, distribution and utilisation of metabolic reserves in *Culex quinquefasciatus* (Diptera: Culicidae): Implications for disease transmission and vector control. *The Journal of Basic and Applied Zoology*, 79(1), 32. [\[Crossref\]](#)
- Wilke, A. B. B., Chase, C., Vasquez, C., Carvajal, A., Medina, J., Petrie, W. D., & Beier, J. C. (2019). Urbanization creates diverse aquatic habitats for immature mosquitoes in urban areas. *Scientific Reports*, 9(1), 15335. [\[Crossref\]](#)
- Wohl, M., & McMeniman, C. J. (2022). Batch rearing *Aedes aegypti*. *Cold Spring Harbor Protocols*. [\[Crossref\]](#)
- World Health Organization. (2022). *World malaria report 2022*. WHO
- World Health Organization. (2024). *World malaria report 2024: Addressing inequity in the global malaria response*. WHO
- Zengenene, M. P., Munnenga, G., Okumu, F. O., & Koekemoer, L. L. (2022). Effect of larval density and additional anchoring surface on the life-history traits of a laboratory colonized *Anopheles funestus* strain. *Medical and Veterinary Entomology*, 36(2), 168-175. [\[Crossref\]](#)
- Zogo, B., Ahoua Alou, L. P., Koffi, A. A., Fournet, F., Dahounto, A., Dabiré, R. K., Baba-Moussa, L., Morou, N., & Penner, C. (2019). Identification and characterization of *Anopheles* spp. breeding habitats in the Korhogo area in northern Côte d'Ivoire: A study prior to a *Bti*-based larvicide intervention. *Parasites & Vectors*, 12(1), 146. [\[Crossref\]](#)