

## ORIGINAL RESEARCH ARTICLE

## Antibacterial Activities of Three (3) Selected Medicinal Plants against Diarrhea Causing Pathogens.

Murtala Saadu Zamfarawa<sup>1\*</sup>, Nasiru Usman Adabara<sup>2</sup>, Faruk Adamu Kuta<sup>3</sup>, Hadiza Lami Muhammed<sup>4</sup>  
Mohamed Wuna<sup>5</sup> , Sa'adu Usman<sup>6</sup> and Isah Musa Gummi<sup>7</sup> .

<sup>1</sup>Department of Microbiology, Faculty of Life Sciences, Aliero, Nigeria.

<sup>2,3,4,5</sup>Department of Microbiology, Faculty of Life Sciences, Federal University of Technology, Minna, Nigeria.

<sup>6</sup>Department of Microbiology Bayero University Kano

<sup>7</sup>Department of Microbiology, Kebbi State University of Science and Technology, Aliero

### ABSTRACT

Diarrhoea is one of the leading causes of morbidity and mortality worldwide, especially in developing countries in sub-Saharan Africa and Southeast Asia. Extracts from *Acacia nilotica* (*Bagaruma*), *Carica papaya* (Gwanda) and *Khaya senegalensis* (Marke) were prepared using the cold maceration method. Well diffusion method was used on Muller Hinton agar and thiosulfate citrate bile salt agar (TCBS), and suspension of the test organisms was prepared in peptone water to match 0.5 McFarland standard, each extract solution at 200, 250 and 300mg/ml, was introduced into respective wells, incubated at 37°C for 24 hours. Ciprofloxacin, at 8 mg/ml, was used as a control. Plant crude extracts of *C. papaya* yielded (19.05g), *A. nilotica* (18.67g) and *K. senegalensis* (13.94g), respectively. The extracts exhibited varying degrees of inhibition against *V. cholerae*, *K. pneumoniae* and *S. enterica* serovar. *Khaya senegalensis* had the highest significant antibacterial activity at 300mg/ml (36.00±0.81f, 19.00±0.98, 22.00±0.69) and was the most active among the three plant extracts, followed by *C. papaya* at 300mg/ml (29.33±1.17, 31.50±0.29, 9.00±1.15) then *A. nilotica* at 200mg/ml, 29.00±0.87, 12.00±0.58, 20.00±0.29. Only *C. papaya* showed no activity at all concentration against *V. cholera*. The MIC of the plant crude extracts against *Vibrio cholerae*, *Klebsiella pneumoniae* and *Salmonella enterica* serovar *Kentucky* was 0.96±0.04, 0.96±0.05 and 1.80±0.17mg/mL, respectively, while MBC was 1.02±0.12, 1.04±0.29 and 1.92±0.23 mg/mL. In conclusion, the three plant crude extracts of (*K. senegalensis*, *A. nilotica* and *C. papaya*) showed varying degrees of antimicrobial activities against the test organisms. Therefore, the leaves of the plants *Carica papaya*, *Acacia nilotica* and *K. senegalensis* can be recommended for diarrhoeal therapy in accordance with the traditional uses.

### ARTICLE HISTORY

Received July 21, 2023.

Accepted September 30, 2023.

Published June 23, 2024.

### KEYWORDS

Diarrhea Pathogens, plant extracts, Antibacterial activity.

© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

## INTRODUCTION

In developing countries, especially in Sub-Saharan Africa, pathogenic diarrhoea is one of the leading causes of morbidity and mortality (Cohen *et al.*, 2022). Diarrhoea-associated bacterial pathogens include *Escherichia coli*, *Vibrio cholera*, Species of *Shigella*, *Salmonella*, *Proteus*, *Yersinia*, and *Campylobacter* (WHO, 2015). Most diarrhoea-associated morbidities and mortalities occur in low to medium-income countries, usually in rural areas. The incidence is fuelled by the vicious cycles of poverty, ignorance, malnutrition, and endemic infections (Mohammed *et al.*, 2014). Herbal preparations are an important traditional medicine therapy that were still used by approximately 80% of the world's population, particularly in developing countries (Kamboj, 2000). A

large number of plant species have been documented for the treatment of various ailments and serve as remedies for human disease (Ibrahim *et al.*, 2016). Furthermore, research has consistently shown that most compounds found in plants have beneficial qualities with little adverse effects, and their derivatives have several medicinal and other desirable properties (Farouk *et al.*, 2008).

In Nigeria, diarrhoeal diseases are the third leading cause of death among the populace, with children under five years more affected (WHO, 2015). In 2017, about 1.7 million people died from diarrheal diseases globally (Akpan *et al.*, 2012). Conventional diarrhoeal antibiotics are not quite affordable, and adverse effects that accompany their intake cannot be ignored (Ramos *et al.*,

**Correspondence:** Murtala Saadu Zamfarawa. Department of Microbiology, Faculty of Life Sciences, Aliero, Nigeria. ✉ [murtala123@gmail.com](mailto:murtala123@gmail.com). Phone Number: +234 703 373 9861.

**How to cite:** Zamfarawa, M. S., Usman, N. A., Adamu, F. K., Muhammed, H. L., Wuna, M., Usman, S., & Musa, I. G. (2024). Antibacterial Activities of Three (3) Selected Medicinal Plants against Diarrhea Causing Pathogens. *UMYU Scientifica*, 3(2), 127 – 132. <https://doi.org/10.56919/usci.2432.013>

2008) Spread of drug-resistant microbes remains a major challenge to healthcare providers, particularly in developing nations like Nigeria (Seddik *et al.*, 2005), and most of the plants used have limited potentials of the most bacteria causing diarrhoea, the need to search for an alternative antimicrobial agent of plant origin cannot be overemphasized. Therefore, this research aims to determine the antibacterial Activities of Three (3) Selected Medicinal Plants against Diarrhea Causing Pathogens.

## MATERIALS AND METHODS

### Study Area

The study was carried out in General Hospital Minna, Niger state, Nigeria, which is the major hospital attended by the populace. Minna is a city in Middle Belt Nigeria, consisting of two Major ethnic groups: Nupe and Gwari. Minna Lies between Latitude 9.58360 N and Longitude 6.54630 E at an altitude of 256m above sea level and has a land area of about 88 km<sup>2</sup> (<http://www.minna.climatetemps.com/map.php>).

### Sample collection

A stool sample from diarrhoea patients was collected in sterile sample bottles, packaged in icepacks and transported to the Center for Genetic Engineering and Biotechnology Laboratory, Federal University of Technology, Minna.

### Isolation of diarrhoea-causing pathogens

Samples collected were inoculated onto MacConkey agar, Eosine methylene blue (EMB) agar and Thiosulfate citrate bile-salts sucrose (TCBS) agar and incubated at 37°C for 24 hours. Distinct colonies were sub-cultured repeatedly to obtain pure cultures of diarrhoea-causing pathogens, which were stored on nutrient agar slant for further use.

### Identification of isolated pathogens

Suspected bacterial diarrhoea-causing pathogens obtained were identified using colony morphology and conventional biochemical tests, including Gram staining, Oxidase test, Voges proskauer test, Indole test, Methyl red, Citrate test, Catalase, Urease and Motility test.

### Gram staining

A thin smear of the pure 24-hour-old culture was prepared on a clean, grease-free slide; it was allowed to dry and fixed by passing over a gentle flame. The smear was then stained by adding 2 drops of crystal violet (Primary stain) solution for 60 seconds and then rinsed with water. The smear was again flooded with Lugol's iodine (Mordant) for 30 seconds and rinsed with water before decolourizing with 70% alcohol. The smear was then counter-stained with Safranin for 60 seconds before rinsing with water and then air drying. The smear was mounted on a microscope and observed under an oil immersion objective lens (x100). Gram-negative cells appeared pink or red, while gram-positive organisms appeared purple.

### Plant Sample collection and identification

Fresh leaves of five medicinal *Carica papaya* (KSUSTA/PSB/H/109C), *Khaya senegalenses* (KSUSTA/PSB/H/61A) *Acacia nilotica* Wild. (KSUSTA/PSB/H/284 was collected in Minna, Niger state. The plants were identified at the Department of Plant Biology School of Life Sciences, Kebbi State University of Science and Technology Aliero Kebbi State.

### Extraction of the crude extracts

The plant extraction procedure was carried out according to the method described by AOAC (2010). The plant leaves were dried under shade at room temperature for at least 7 days, segregated and pulverized by a mechanical grinder, forming a coarse powder. 100g of powdered samples were weighed and macerated in 500ml of n-hexane, ethyl acetate and methanol for 72 h in a 1:5 (w/v) ratio, respectively. The supernatants obtained were filtered using Whatman No 1 filter paper and evaporated until dried under reduced pressure (204 mbar) at 40 °C.

### Antimicrobial assay

The antibacterial activity of the selected plant extracts was tested using the agar well diffusion technique on Muller Hinton agar according to the method described by Agarry *et al.* (2005)\*. Suspension of the test organisms was prepared in peptone water to match turbidity of 0.5 Macfarland standard and inoculated onto Mueller Hinton agar. Agar wells with a diameter of 8mm were punched aseptically using a sterile cork borer. Each extract solution at the desired concentration (50, 70 and 100mg/ml) was introduced into each well and then incubated at 37°C for 24 hours. The extracts' antimicrobial activity was measured according to the National Committee for Clinical Laboratory Standards (NCCLS, 2002)\*.

### Minimum Inhibitory Concentration (MIC) and Maximum Bactericidal Concentration (MBC)

Based on the results of the antibacterial testing, the most efficient extracts were chosen to identify its MIC and MBC. The Broth tube dilution method was used as described by Srikacha and Ratananikom, (2020)\*. Serial dilution was adopted to prepare various concentrations of the selected extracts in test tubes, which were inoculated with a standardized number of organisms and incubated at 37°C for 18-24 hours. The lowest concentration (highest dilution) of the extract with no visible growth of the test organism was considered the MIC. The MBC was determined by plating the contents of the MIC tubes onto nutrient agar plates. The lowest extract concentration with minimum growth of the test organism on the agar plate was considered the MBC.

## RESULTS

### Extraction Yields of Plants

The extraction yield of a plant used in this study is shown in Table 2: *Carica papaya* (19.05g), *Acacia nilotica* (18.67g) and the lowest yield was observed in *Khaya senegalensis* (13.94g).

**Table 1: Identity of test organisms**

Sample ID	Isolate code	Scientific Name	E value	Per. Ident	Accession
1	KP	<i>Klebsiella pneumoniae</i> strain IAUK 8738	9e-171	98.57%	MK571203.1
2	ST	<i>Salmonella enterica</i> subsp. <i>enterica</i> serovar <i>Kentucky</i> strain Medellin Colombian	0	97.40%	MH445517.1
3	VC	<i>Vibrio cholerae</i> strain RP01	0	100.00%	MH174981.1

**Table 2: Yield of Extract from Plants used in this study**

Plants	Extraction Yield (g)
<i>Acacia nilotica</i>	18.67
<i>Carica papaya</i>	19.05
<i>Khaya senegalensis</i>	13.94

Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of Crude Extracts  
The MIC and MBC of the crude extracts of *Acacia nilotica* and *Khaya senegalensis* are shown in Table 3. The Minimum

Inhibitory Concentration (MIC) of *Acacia nilotica* against *Vibrio cholera*, *Klebsiella pneumoniae* and *Salmonella enterica* serovar *Kentucky* was observed at  $0.96 \pm 0.03$ ,  $0.19 \pm 0.00$  and  $0.96 \pm 0.06$  mg/mL respectively, while the minimum bactericidal concentration (MBC) was observed at  $24.00 \pm 0.87$ ,  $0.96 \pm 0.05$  and  $4.80 \pm 0.17$  mg/mL respectively. For *Khaya senegalensis*, MIC was observed at  $0.96 \pm 0.04$ ,  $0.96 \pm 0.05$  and  $4.80 \pm 0.17$  while MBC was recorded at  $4.80 \pm 0.12$ ,  $4.80 \pm 0.29$  and  $4.80 \pm 0.23$ . (0.64g) followed by aqueous (0.57g) and then ethyl-acetate (0.25g). Similarly, n-hexane had the highest yield (0.46g) in *Khaya senegalensis*, followed by ethyl-acetate (0.28g) and then aqueous (0.25g).

**Table 3: Minimum Inhibitory and Maximum Bactericidal Concentrations**

Plant Extract	MIC			MBC		
	<i>V. cholerae</i>	<i>K. pneumoniae</i>	<i>S. enterica</i> serovar <i>Kentucky</i>	<i>V. cholerae</i>	<i>K. pneumoniae</i>	<i>S. enterica</i> serovar <i>Kentucky</i>
<i>C. papaya</i>	$0.96 \pm 0.03$	$0.19 \pm 0.00$	$0.96 \pm 0.06$	$1.09 \pm 0.87$	$0.20 \pm 0.05$	$1.04 \pm 0.17$
<i>K. senegalensis</i>	$0.96 \pm 0.04$	$0.96 \pm 0.05$	$1.80 \pm 0.17$	$1.02 \pm 0.12$	$1.04 \pm 0.29$	$1.92 \pm 0.23$
Control	0.005	0.005	0.020	0.005	0.005	0.020

Values are presented as mean  $\pm$  standard error of triplicates' mean (SEM).

MIC = Minimum Inhibitory; MBC = Minimum Bactericidal Concentrations

Antibacterial Activity of Crude Plant Extracts against Test Organisms

and *Salmonella enterica* serovar *Kentucky* is shown in Table 4 below.

The Antibacterial activity of three different plant extracts used in this study against *Vibrio cholera*, *Klebsiella pneumoniae*

**Table 4: Inhibitory Effects of Methanolic Crude Extracts Against Bacteria (mm).**

Extract	Concentration (mg/mL)	<i>Vibrio cholerae</i>	<i>Klebsiella pneumoniae</i>	<i>Salmonella enterica</i> serovar <i>Kentucky</i>
<i>Acacia nilotica</i>	200	$11.00 \pm 0.58^b$	$22.00 \pm 0.87^e$	$14.00 \pm 0.58^c$
	250	$25.33 \pm 0.60^c$	$31.50 \pm 0.87^f$	$11.00 \pm 0.58^{ab}$
	300	$29.33 \pm 1.17^d$	$31.50 \pm 0.29^f$	$9.00 \pm 1.15^a$
<i>Carica papaya</i>	200	$0.00 \pm 0.00^a$	$9.00 \pm 1.15^a$	$18.00 \pm 0.58^{ef}$
	250	$0.00 \pm 0.00^a$	$17.00 \pm 1.15^d$	$26.33 \pm 0.60^h$
	300	$0.00 \pm 0.00^a$	$24.00 \pm 0.58$	$28.33 \pm 1.17^h$
<i>Khaya senegalensis</i>	200	$29.00 \pm 0.87^d$	$12.00 \pm 0.58^b$	$20.00 \pm 0.29^g$
	250	$32.50 \pm 0.29^e$	$16.20 \pm 1.27^{cd}$	$20.00 \pm 1.15^g$
	300	$36.00 \pm 0.81^f$	$19.00 \pm 0.98^d$	$22.00 \pm 0.69^g$
Control	8mg/ml	$24.21 \pm 0.82$	$12.03 \pm 13$	$32.07 \pm 1.43$

Keys: Values are presented as mean  $\pm$  standard error of the mean (SEM) of triplicates

Values with different superscripts in a column are significantly different at  $p < 0.05$

## DISCUSSION

## Antibacterial Activity of Crude Plant Extracts against Test Organisms

*Carica papaya*, *Acacia nolitica* and *Khaya senegalensis* are well-known medicinal plants with health benefits against many ailments (Doughari *et al.*, 2007; Ahmad and Wudil, 2013\*; Ibrahim *et al.*, 2013; Galalain *et al.*, 2019; Namadina *et al.*, (2021). Extracts from these plants used in this study exhibited varying degrees of antibacterial activity against *Vibrio cholera*, *Klebsiella pneumoniae* and *Salmonella enterica* serovar *Kentucky*. *Carica papaya* and *Khaya senegalensis* were noted to be the most active among the three plant extracts tested, as they showed significantly higher zones of inhibition at all concentrations against all the test organisms.

The efficacy of the plant extracts evaluated was concentration-dependent, as the antimicrobial activity increases with increasing concentration of the plant extracts, indicated by the increasing diameter of each inhibition zone. The observed inhibitory activity of the plant extracts against these diarrhoea-causing pathogens validates the use of the plant for treating diarrhoea in traditional medicine. This result is consistent with findings reported by Doughari *et al.* (2007), Mann *et al.* (2014), Kuta *et al.* (2015), Abdallah *et al.* (2016), Ali *et al.* (2017) Sadiku *et al.* (2020) with few exceptions. They all reported varying ranges of inhibitory effects of these plant extracts against various bacterial pathogens at various concentrations. According to Kuta *et al.* (2015), leaf and stem extracts of *Khaya senegalensis* showed inhibitory activity against *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Streptococcus pneumoniae* and *Escherichia coli* at concentrations ranging from 400 to 1000mg/ml,

Similarly, Abdallah *et al.* (2016) reported leaf and stem extracts of *Khaya senegalensis* to have inhibitory activity against diarrheal/stool isolates (*E. coli*, *Shigella* sp. and *Salmonella* sp.). Musah (2019) observed that matured leaf extract of *Polioctigma thoningii* showed antibacterial activity against *E. coli* and *S. enterica* Typhi at 10 and 5mg/cm<sup>3</sup> concentrations but had no activity against *P. aeruginosa*. From the study of Evbuomwan *et al.* (2018)\*, *Vernonia amygdalina* leaf extract showed significant antibacterial activity against multidrug-resistant bacterial isolated, namely *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Klebsiella pneumoniae* at various concentrations ranging from 25 to 200mg/mL.

Mann *et al.* (2014) observed that stem bark extract of *Anogeissus leiocarpus* showed an inhibitory effect against multidrug-resistant *Staphylococcus aureus* and food pathogen of *Hibiscus sabdariffa calxy* (Zobo) drink, respectively. The antimicrobial activity of these plant extracts is usually attributed to their phytochemical constituent. Although the presence and roles of phytochemicals were not investigated in this study, previous studies have shown Tannins, Flavonoids, Saponins and Alkaloids, which are usually reported to be present in these plants, to have

antibacterial properties (Mann *et al.*, 2014; Abalaka *et al.*, 2016;).

Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration of Crude Extracts of *Carica papaya* and *Khaya senegalensis* against test isolates

The minimum inhibitory concentration (MIC) is the smallest concentration that visibly inhibits growth. The MIC is useful in determining the smallest effective dosage of a substance against an organism (Kuta *et al.*, 2015). The MIC of *Carica papaya* against *Vibrio cholera*, *Klebsiella pneumoniae* and *Salmonella enterica* serovar *Kentucky* was observed at 0.96±0.03, 0.19±0.00 and 0.96±0.06mg/mL, respectively, while the minimum bactericidal concentration (MBC) was observed at 1.09±0.04, 0.20±0.05 and 1.04±0.17mg/mL respectively. For *Khaya senegalensis*, MIC was detected at 0.96±0.04, 0.96±0.05 and 1.80±0.17 mg/mL concentration, while MBC was recorded at 1.02±0.12, 1.04±0.29 and 1.92±0.23 mg/mL concentration. Plant extracts are considered to have a good inhibitory activity if they present MIC value ≤ 100mg/mL, a moderate inhibitory activity when MIC ranges between 100 to 500mg/mL, a weak inhibitory activity if MIC value ranges between 500 to 1000mg/mL and no inhibitory activity when > 1000mg/mL (Jafari *et al.*, 2008; Mann *et al.*, 2014).

Considering this report, the MIC and MBC values recorded from the antibacterial activity of the present study showed good inhibitory activity. The current findings lend credence to the traditional use of this plant as a medicine for infectious diseases, particularly those caused by the test organisms susceptible to the extracts. However, this result is contrary to the findings of Kuta *et al.* (2015), Abdallah *et al.* (2016), Ali *et al.* (2017), Usman *et al.* (2020) and Amadou *et al.* (2020) who all reported much higher MIC and MBC values against various bacterial pathogens. Ugoh *et al.* (2014) observed methanolic and ethanolic stem bark extracts of *Khaya senegalensis* with a MIC value of 250 and 200mg/mL against *Salmonella enterica* subsp. *enterica* serovar Typhi. MIC value of 25mg/ml was reported by Shinde *et al.* (2020) for methanolic stem bark extract of *Khaya senegalensis* against both *Klebsiella pneumoniae* and a carbapenem-resistant strain of *Escherichia coli*.

In Kuta *et al.* (2015) studies, the aqueous and ethanolic leaf extract of *Khaya senegalensis* had MIC values of 400 and 200mg/mL, respectively, against *Pseudomonas aeruginosa*, *Streptococcus pneumoniae* and *Escherichia coli*. The study also reported MBC values of 800 and 400mg/mL for aqueous and ethanolic leaf extract of *Khaya senegalensis* against the same organisms, respectively. Ali *et al.* (2017) studied the antimicrobial activity of methanolic stems bark extracts of *Anogeissus leiocarpus* against several bacterial isolates; the study reported MIC and MBC values of 10 and 40 mg/mL for *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Salmonella typhi*, *Streptococcus faecalis* and *Corynebacterium ulcerans*. In Chomini *et al.* (2021)

For the aqueous extract, MIC was detected at 100, 200 and 100 mg/mL, while the ethanolic extract had MIC values at 200, 400 and 200mg/mL against *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Streptococcus mutans* respectively. While the MBC was detected at 400mg/mL for both aqueous and ethanolic extract against all the isolates. The variation in the MIC and MBC values reported in these studies could be due to the phytochemical composition of their respective extracts, which is usually dependent on the polarity of the solvents used for extraction, as solvent type tends to influence the kind of bioactive compound released from the plant materials. Also, variation in results could be attributed to the genetic makeup of each test organism used. Different organisms have been shown to respond differently to different and same concentrations of a specific antimicrobial substance (Sadiku *et al.*, 2020).

## CONCLUSION

Various diarrhoea-causing pathogens were isolated from stool samples of patients attending General Hospital, Minna. They include *Salmonella enterica* serovar *Kentucky* strain Medellin Colombian, *Klebsiella pneumoniae* strain IAUK 8738 and *Vibrio cholerae* strain RP01. Crude extract of the three (3) plants showed varying degrees of antimicrobial activity against the test organisms, with *Khaya senegalensis*, *Acacia nitolytica* and *Carica papaya* being the most active.

## REFERENCES

Abalaka M. E., Akpor O. B., Bello A. B., Olonitola O. S., Sani A. and Osemwegei O. O. (2016). Synthesis of Silver Nano Particles (AgNPs) from Bark and Rooth of African Mahogany (*Khaya senegalensis*) and the Comparative Study of their Antimicrobial Properties. *Asian Journal of Microbiology and Biotechnology* 1(1): 25-32. [\[Crossref\]](#)

Abdallah, M. S., Ahmed, I., & Yahaya, S. M. (2016). Antibacterial activity and phytochemical screening of leaf and stem (bark) extract of *Khaya senegalensis* against Diarrhoeal stool isolates. *Cibtech Journal of Microbiology*, 5(1), 17-24.

Agarry O.O, Olaleye M.T and Bello-Michael, C. O. (2005) Comparative antimicrobial activities of Aloe vera gel and leaf. *African Journal of Biotechnology* Vol. 4 (12), pp. 1413-1414,

Ahmad I.M. and Wudil A.M. (2013). Phytochemical Screening and Toxicological Studies of Aqueous Stem Bark Extract of *Anogeissus leiocarpus* in Rats. *Asian Journal of Scientific Research* 6 (4) 781-788.

Akpan, M. M., Odeomena, C. S., Nwachukwu, C. N. and Danladi, B. (2012). Antimicrobial assessment of ethanolic extract of *Costus afer* Leaves. *Asian Journal of Plant Science and Research*, 2(3), 335-341

Ali, S. A. M., Elbadwi, S. M. A., Idris, T. M., and Osman, K. M. (2011). Hepatoprotective activity of

aqueous extract of *Khaya senegalensis* bark in rats. *Journal of Medicinal Plants Research*, 5(24), 5863-5866.

Ali, S. A. M., Elbadwi, S. M. A., Idris, T. M., and Osman, K. M. (2017). Hepatoprotective activity of aqueous extract of *Khaya senegalensis* bark in rats. *Journal of Medicinal Plants Research*, 5(24), 5863-5866.

Amadou, I., Soulé, M., and Salé, A. (2020). An overview on the importance of *Acacia nilotica* (L.) willd. ex del.: A review. *Asian Journal of Research in Agriculture and Forestry*, 5, 12-18. [\[Crossref\]](#)

AOAC Association of Official Analytical Chemists, AOAC (2010). *Official methods of analysis. Association of Official Analytical Chemists*, 15th Edition Washington D.C., USA. 184pp

Chomini, S. M., Joshua, V. I., Ukwadi, M., Sadiku, Y., Popoola, S. A., Imoh, J. A., and Usman, S. A. (2021) Phytochemical Screening and Antibacterial Activities of *Anogeissus leiocarpa* (DC.) Gull. and Perr. Leaf Extracts against *Escherichia coli* (T. Escherichia) and *Salmonella typhimurium* (Loeffler). *Australian Journal of Science and Technology*, 5(2), 527-532.

Cohen, A. L., Platts-Mills, J. A., Nakamura, T., Operario, D. J., Antoni, S., Mwenda, J. M. & Serhan, F. (2022). Aetiology and incidence of diarrhoea requiring hospitalisation in children under 5 years of age in 28 low-income and middle-income countries: findings from the Global Pediatric Diarrhea Surveillance network. *BMJ Global Health*, 7(9), e009548. [\[Crossref\]](#)

Doughari, J. H. (2012). *Phytochemicals: extraction methods, basic structures and mode of action as potential chemotherapeutic agents* (pp. 1-33). Rijeka, Croatia: INTECH Open Access Publisher.

Evbuomwan, I; Chukwuka, Ep; Obazenu, Ei; Ilevbare, L (2018) Antibacterial Activity of *Vernonia amygdalina* Leaf Extracts against Multidrug Resistant Bacterial Isolates. *J. Appl. Sci. Environ. Manage* 22 (1) 17-21. [\[Crossref\]](#)

Farouk, M.A, Musa, A., and Yakasai I.A. (2016). Determination of Cadmium and Lead Levels In Meat, Liver And Kidney Of Cow And Goat Marketed In Zaria, Nigeria. *Bayero Journal of Pure and Applied Sciences*, 9(1) 180 – 184. [\[Crossref\]](#)

Galalain, M. A., Habib, A. A., Tukur, S., Bashir, A. R., and Abdu, K. (2019). Antifungal efficacy of *Anogeissus leiocarpus* DC against *Aspergillus niger* and *Rhizoctonia spp.* *Fudma Journal of Sciences-ISSN: 2616-1370*, 3(2), 179-184.

Ibrahim, M. A., Musa, A. M., Aliyu, A. B., Mayaki, H. S., Gideon, A., and Islam, M. S. (2013). Phenolics-rich fraction of *Khaya senegalensis* stem bark: antitrypanosomal activity and amelioration of

- some parasite-induced pathological changes. *Pharmaceutical Biology*, **51**(7), 906-913. [\[Crossref\]](#)
- Jafari, F., Hamidian, M., Rezadehbashi, M., Doyle, M., Salmanzadeh-ahrabi, S., Derakhshan, F. & Zali, M. R. (2008). Prevalence and antimicrobial resistance of diarrheagenic *Escherichia coli* and *Shigella* species associated with acute diarrhea in Tehran, Iran. *Canadian Journal of Infectious diseases and Medical Microbiology*, **19**(3), e56-e62. [\[Crossref\]](#)
- Kamboj, V. P. (2000). Herbal medicine. *Current science*, **78**(1), 35-39.
- Kuta F. A., Tsado D. J., Garba S. A. and Saidu N. A. (2015). Antibacterial Activity of the Leaf and Stem bark crude extract of *Khaya senegalensis*. *Medicinal and Aromatic Plant Research Journal* Vol. **3**(1) pp 9-15
- Mann, A., Yusuf, A. and Daniyan, S. (2014). TLC analysis and bioactivity screening of the stem bark extract of *Anogeissus leiocarpus* against multi-resistant *Staphylococcus aureus* and quantification of its phytoconstituents. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, **5**, (187-203).
- Mohammed A., K.Ghoneim, M. Amer, A. Al-Daly, F. Khadrawy and M.A. Mahmoud (2014). Transaminase perturbation in certain tissues of *Schistocerca gregaria*(Forsk.) (Orthoptera: Acrididae) by *Punica granatum* Linn. (Lythraceae) extracts
- Mohammed, S. and Tamiru, D. (2014). The burden of diarrheal diseases among children under five years of age in Arba Minch district, Southern Ethiopia, and associated risk factors: A cross-sectional study. *International Scholarly Research Notices* 2014, 654901. [\[Crossref\]](#)
- Muhammad, B., Sadiq, I., Joel, T., Tay, Z., Aye, C. and Anil, K.A. (2015). Antibacterial Activities and Possible Modes of Action of *Acacia nilotica* (L.) Del. against Multidrug-Resistant *Escherichia coli* and *Salmonella*. [\[Crossref\]](#)
- Musah, S. (2019). Antibacterial activity of matured leaf extract of *Poliostigma thoningii* against *E. coli* and *S. enterica* Typhi at various concentrations. *African Journal of Biotechnology*, **18**(12), 399-406. [\[Crossref\]](#)
- Namadina, M. M., Mukhtar, A. U., Karaye, S. I., Musa, F. M., Bah, I. H., Maitama, F. Y. (2021) Biological Activity and Phytochemical Constituents of *Anogeissus leiocarpus* Stem Bark Extracts. *Bayero Journal of Pure and Applied Sciences*, **14**(1), 85- 94, [\[Crossref\]](#)
- NCCLS (2002) Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals, Approved Standard. 2nd Edition, NCCLS Document M31-A2 22(6), Clinical and Laboratory Standards Institute, Wayne. [\[Crossref\]](#)
- Nikom Srikacha and Khakhanang Ratananikom (2020). Antibacterial activity of plant extracts in different solvents against pathogenic bacteria: An in vitro experiment. *Journal of Acute Disease.* ; **9**(5): 223-226. [\[Crossref\]](#)
- Ramos Morales, E. ; Molina Alcaide, E. ; Sanz Sampelayo, M. R. S., (2008). Milk production of dairy goats fed diets with different legume seeds: Effects of amino acid composition of the rumen undegradable protein fraction. *J. Sci. Food Agric.*, **88** (13): 2340-2349. [\[Crossref\]](#)
- Seddik S. Bun , H. Bun , D. Guédon, C. Rosier , E. Ollivier (2005). Effect of green tea extracts on liver functions in Wistar rats *Journal of Food and Chemical Toxicology*. Volume 44, Issue 7, July 2006, Pages 1108-1113
- Shinde, A., Kolhe, D., Hase, D. P., and Chavan, M. J. (2020). A brief introduction about *Carica papaya* linn. *World Journal of Pharmaceutical Research.* **9**(12), 237-244.
- Ugoh Sylvanus Chukwudi \* Oluwabunmi Olaitan Agarry, and Samuel Alimi Garba (2014). studies on the antibacterial activity of *Khaya senegalensis* [(Desr.) A. Juss] stem bark extract on *Salmonella enterica* subsp. *enterica* serovar Typhi [(ex Kauffmann and Edwards) Le Minor and Popoff]. *Asian Pac J Trop Biomed.* 79-S283. [\[Crossref\]](#)
- Usman, A. H., Ali, T., & Danjani, A. G. (2020). Phytochemical and antimicrobial studies of stem-bark extracts of *Anogeissus leiocarpus* found in Dutsin-ma, katsina-nigeria. *Fudma Journal of Sciences*, **4**(2), 156-167. [\[Crossref\]](#)
- WHO Cholera, vaccines: WHO, position paper – August 2017, *Epidemiological Record* 25 August 2017, No 34, 2017, 92, 477–500.
- World Health Organization . Cholera, (2015). *Weekly Epidemiological Record.* 96:445-54.