

ORIGINAL RESEARCH ARTICLE

Commutativity of some Prime Narrings using Left-Sided outer (σ, τ) -n-Derivation

Moharram Ali Khan¹ , Ibrahim Shamsu² .

^{1,2}Department of Mathematics and Statistics, Umaru Musa Yar'adua University, Katsina, Nigeria..

ABSTRACT

In the field of mathematics, pure mathematics is very important as it gives rise to the basis for the formation of all applicable mathematical concepts in solving real-life problems. Algebra is such an integral part of pure mathematics. It consists of the Ring theory. It has been discovered that there exists some structures similar to rings with little deformity and they are called NEARRINGS. These structures do not commute mostly as they fail to satisfy distributive law. To ascertain the commutativity of nearrings, we need derivation(s). Because several papers had been presented dealing with left-nearrings, this paper aimed to consider right-nearrings which has not been done before in that respect, to the best of our knowledge. Some methods dealing with left-nearrings have been studied and modified in this work, and new derivation has been introduced to take care of right-nearrings. For Let n be a positive integer, N be a right nearring, and automorphisms $\sigma, \tau: N \rightarrow N$, in this study, the concept of left-sided outer (σ, τ) - n -derivations on the nearrings is introduced, and several features are examined to show commutativity of prime right nearring using the derivation. The results obtained in this paper show that the right-nearrings are commutative when subjected to derivation.

ARTICLE HISTORY

Received August 14, 2023.
Accepted November 22, 2023.
Published December 21, 2023.

KEYWORDS

Commutativity, Derivations, Nearrings, Outer (σ, τ) - n -derivation, Prime nearrings.

© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

INTRODUCTION

An algebraic structure or a triple $(N, +, \cdot)$ containing a non-null set N together with binary operations (addition '+' as well as multiplication ' \cdot ') that meet all the axioms of rings except for commutativity of addition and just one distributive law holds, is called a nearring. Precisely, right nearring is a triple $(N, +, \cdot)$ as such, (i) $(N, +)$ makes a group (ii) (N, \cdot) is a semi-group (iii) $(s + t)r = sr + tr$, for all $r, s, t \in N$. See Clay (1992) and Pilz (1983) to learn more.

In this article, the right nearring is denoted by N and τ are automorphisms of N . The nearrings' centre is indicated by Z and is defined to be a collection $\{x \in N | xy = yx, \forall y \in N\}$. A right near-ring N is called a prime nearring if $xNy = \{0\}$ then $x = 0$ or $y = 0$, for all $x, y \in N$. N is semiprime nearring if, $xNx = \{0\}$ then $x = 0$ for all $x \in N$ and N is commutative if (N, \cdot) is commutative (that is $xy - yx = 0 \forall x, y \in N$). The symbol $[x, y]$ will be denoted by $xy - yx$ for $x, y \in N$ and is called Lie product (multiplicative commutator) of x and y .

Prime nearrings: these are a special class of nearrings with the property that $xNy = \{0\}$ then $x = 0$ or $y = 0$, for

all $x, y \in N$, where N is the nearring of reference. These are very important, especially in the study of commutativity, as they enable us to use the primeness property to test the commutativity stance of the structure. Some examples of prime nearrings include: [1] Let $(N, +)$ be a group; we define an operation $*$ on N by $x * y = x$ for all $x, y \in N$, it is observed that $(N, +, *)$ is a right prime nearring. [2] Let $(N, +)$ be a Klen four group. Consider the following table;

*	0	r	s	t
0	0	0	0	0
r	0	0	r	r
s	0	r	s	s
t	0	r	t	t

Clearly $(N, +, *)$ is prime right nearring, since for $r, s \in N \Rightarrow r * N * s = \{0\}$ only if $r = 0$ or $s = 0$.

Correspondence: Ibrahim Shamsu. Department of Mathematics and Statistics, Umaru Musa Yar'adua University, Katsina, Nigeria. ✉ shamsuibrahim0043@gmail.com. Phone Number: +234 809 697 8230.

How to cite: Moharram A. K. & Ibrahim S. (2023). Commutativity of some Prime Nearrings using Left-Sided outer (σ, τ) - n -Derivation. *UMYU Scientifica*, 2(4), 108 – 113. <https://doi.org/10.56919/usci.2324.013>

Also; $\frac{\delta(P_1 P_2)}{P_2 + 0 \cdot \overline{P_1}} = \frac{\sigma(P_1)\delta(P_2) + \tau(P_2)\delta(P_1)}{\overline{P_1} \cdot \overline{P_2}} = \overline{P_1} \cdot \overline{P_2}$.

Generally; $\delta(P_1, \dots, P_n) = \overline{P_i}$ where $i = 1, 2, 3, \dots, n$ and $\overline{P_i} = a_i x$, also $\sigma(P_i) = \overline{P_i}$ and $\tau(P) = 0$, $P_i, \overline{P_i}, P \in N$, then δ is the left-sided outer (σ, τ) - n -derivation on N .

The above is the left-sided outer (σ, τ) - n -derivation on N , which could be used to establish commutativity of special class of right nearrings.

RESULTS

Theorem 1: Assume a prime nearring N and δ be a positive left-sided outer (σ, τ) - n -derivation of N . If $\delta(N, \dots, N)$ is contained in centre Z , it implies N is commutative.

Theorem 2: Let δ be non-zero left-sided outer (σ, τ) - n -derivation on the nearring, assume that for any positive integer i with values $1, 2, 3, \dots, n$. $\delta(x_1, x_2, \dots, [x_i, y_i], \dots, x_n) = 0 \ \forall x_1, x_2, x_3, \dots, x_n \in N$. Implies (N, \cdot) is commutative.

DISCUSSIONS

Proofs of Main Results

We establish the following In order to establish our key theorems;

Lemma 1; Let $\delta: \underbrace{N \times N \times \dots \times N}_{n\text{-times}} \rightarrow N$ an n -additive map. It follows that δ is a left-sided outer (σ, τ) - n -derivation in N if;

$$\begin{aligned} &\sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) \\ &\quad + \tau(y_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \\ &= \tau(y_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \\ &\quad + \sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) \end{aligned}$$

For all $x_i, y_i \in N, i$ is $1, 2, 3, \dots, n$.

Proof;

Let δ be left-sided outer (σ, τ) - n -derivation, and $\tau(y_i)$ is a distributive element and $x_i, y_i \in N$ where $1 \leq i \leq n$, then;

$$\begin{aligned} &\delta(x_1, \dots, (x_i + x_i)y_i, \dots, x_n) \\ &= \sigma(x_i + x_i)\delta(x_1, \dots, y_i, \dots, x_n) \\ &\quad + \tau(y_i)\delta(x_1, \dots, (x_i + x_i), \dots, x_n) \\ &= \sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) + \\ &\quad \sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) + \tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) + \\ &\quad \tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) \end{aligned} \tag{1}$$

Again,

$$\begin{aligned} &\delta(x_1, \dots, x_i y_i + x_i y_i, \dots, x_n) \\ &= \delta(x_1, \dots, x_i y_i, \dots, x_n) + \delta(x_1, \dots, x_i y_i, \dots, x_n) \\ &= \sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) + \\ &\quad \tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) + \sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) \\ &\quad + \tau(y_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \end{aligned} \tag{2}$$

Comparing (1) and (2) yields;

$$\begin{aligned} &\sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) + \tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) \\ &= \tau(y_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \\ &\quad + \sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) \end{aligned}$$

Hence proved.

Lemma 2: let δ be left-sided outer (σ, τ) - n -derivation on N . Then

$$\begin{aligned} &u\{\delta(x_1, \dots, x_i y_i, \dots, x_n)\} \\ &= u\sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) \\ &\quad + u\tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) \end{aligned}$$

For all $x_1, x_2, x_3, \dots, x_i, y_i, \dots, x_n \in N \ 1 \leq i \leq n$.

Proof. Since σ is an automorphism upon $N, \exists v \in N$ as such $\sigma(v) = u \in N$

For all $x_i, y_i \in N \ 1 \leq i \leq n$.

$$\begin{aligned} &\delta(x_1, x_2, x_3, \dots, v(x_i y_i), \dots, x_n) \\ &= \sigma(v)\delta(x_1, x_2, x_3, \dots, x_i y_i, \dots, x_n) \\ &\quad + \tau(x_i y_i)\delta(x_1, x_2, x_3, \dots, v, \dots, x_n) \\ &= \sigma(v)\delta(x_1, x_2, x_3, \dots, x_i y_i, \dots, x_n) + \\ &\quad \tau(x_i)\tau(y_i)\delta(x_1, x_2, x_3, \dots, v, \dots, x_n) \end{aligned} \tag{3}$$

Again;

$$\begin{aligned} &\delta(x_1, x_2, x_3, \dots, (vx_i)y_i, \dots, x_n) \\ &= \sigma(vx_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) \\ &\quad + \tau(y_i)\delta(x_1, x_2, x_3, \dots, vx_i, \dots, x_n) \\ &= \\ &\quad \sigma(v)\sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) + \\ &\quad \tau(y_i)\{\sigma(v)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) + \\ &\quad \tau(x_i)\delta(x_1, x_2, x_3, \dots, v, \dots, x_n)\} \\ &= \\ &\quad \sigma(v)\sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) + \\ &\quad \tau(y_i)\sigma(v)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) + \\ &\quad \tau(y_i)\tau(x_i)\delta(x_1, x_2, x_3, \dots, v, \dots, x_n)\} \\ &= \\ &\quad \sigma(v)\sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) + \\ &\quad \sigma(v)\tau(y_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) + \end{aligned}$$

$$\tau(x_i)\tau(y_i) \delta(x_1, x_2, x_3, \dots, v, \dots, x_n) \} \quad (4)$$

Comparing (3) and (4), we have that

$$\begin{aligned} & \sigma(v)\delta(x_1, x_2, x_3, \dots, x_i y_i, \dots, x_n) \\ &= \sigma(v)\sigma(x_i)\delta(x_1, x_2, x_3, \dots, y_i, \dots, x_n) \\ &+ \sigma(v)\tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) \end{aligned}$$

⇒

$$\begin{aligned} u\delta(x_1, \dots, x_i y_i, \dots, x_n) \\ &= u\sigma(x_i)\delta(x_1, \dots, y_i, \dots, x_n) \\ &+ u\tau(y_i)\delta(x_1, \dots, x_i, \dots, x_n) \end{aligned}$$

Hence proved.

Proof of Theorem 1;

As $\delta(N, N, N, \dots, N) \subseteq Z$, and δ is non-zero, then we say that there exist non-zero elements. $x_1, \dots, x_n \in N$, such that; $\delta(x_1, \dots, x_i, \dots, x_n) \subseteq Z \setminus \{0\}$, i ranges from $1, 2, \dots, n$.

$$\text{Then; } \delta(x_1, \dots, x_i, \dots, x_n) + \delta(x_1, \dots, x_i, \dots, x_n) = \delta(x_1, \dots, x_i + x_i, \dots, x_n) \in Z$$

By Result# (5), then it means N with $+$ is abelian.

We now need to show the commutativity of N .

Let $x, y \in N$. For automorphisms of N are σ and τ , $\exists u, v \in N$ such that $\sigma(u) = x$ and $\tau(v) = y$.

Since $\delta(N, \dots, N) \subseteq Z$, then we assume;

$$\delta(x_1, \dots, x_i, \dots, x_n)y = y\delta(x_1, \dots, x_i, \dots, x_n).$$

Take when $i = 1$ changing x_1 by uv

$$\begin{aligned} & \delta(uv, x_2, x_3, \dots, x_n)y = y\delta(uv, x_2, x_3, \dots, x_n) \\ & \Rightarrow \{\sigma(u)\delta(v, x_2, \dots, x_n) + \tau(v)\delta(u, x_2, \dots, x_n)\}y = \\ & y\{\sigma(u)\delta(v, x_2, \dots, x_n) + \tau(v)\delta(u, x_2, \dots, x_n)\} \\ & \Rightarrow \{x\delta(v, x_2, x_3, \dots, x_n) + y\delta(u, x_2, x_3, \dots, x_n)\}y = \\ & y\{x\delta(v, x_2, x_3, \dots, x_n) + y\delta(u, x_2, x_3, \dots, x_n)\} \end{aligned}$$

By Lemma 2, we have;

$$\begin{aligned} x\delta(v, x_2, x_3, \dots, x_n)y + y\delta(u, x_2, x_3, \dots, x_n)y \\ &= yx\delta(v, x_2, x_3, \dots, x_n) \\ &+ yy\delta(u, x_2, x_3, \dots, x_n) \end{aligned}$$

$$\Rightarrow x\delta(v, x_2, x_3, \dots, x_n)y = yx\delta(v, x_2, x_3, \dots, x_n)$$

Then we have;

$$\delta(v, x_2, x_3, \dots, x_n)(xy - yx) = 0, \text{ for all the elements in } N$$

Then $z\delta(v, x_2, x_3, \dots, x_n)(xy - yx) = 0, \forall z \in N$.

By using the hypothesis and $\delta(N, N, \dots, N) \subseteq Z$,

$$\delta(v, x_2, x_3, \dots, x_n)z(xy - yx) = 0, \forall z \in N$$

$$\delta(v, x_2, x_3, \dots, x_n)N(xy - yx) = \{0\}.$$

As N is a prime, it means $\delta(v, x_2, x_3, \dots, x_n) = 0$ or $(xy - yx) = 0$.

If $xy - yx = 0$, then, the nearring is commutative.

As $\delta(v, x_2, x_3, x_4, \dots, x_n) = 0$.

Substitute v with vx_1 in the equation above and by the hypothesis, we have;

$$x\delta(vx_1, x_2, x_3, \dots, x_n) = \delta(vx_1, x_2, x_3, \dots, x_n)x.$$

By Lemma 2 and the fact that $\delta(N, N, \dots, N) \subseteq Z$, we have;

$$\delta(x_1, x_2, x_3, \dots, x_n)(xy - yx) = 0.$$

Since $\delta(N, N, \dots, N) \subseteq Z$,

$$\delta(x_1, x_2, x_3, \dots, x_n)N(xy - yx) = \{0\}.$$

Since $\delta(x_1, x_2, x_3, \dots, x_n) \neq 0$, $xy - yx = 0 \Rightarrow xy = yx, \forall x, y \in N$.

N is a commutative nearring as a result.

Proof of Theorem 2;

$\forall x_1, x_2, x_3, \dots, x_i, y_i, \dots, x_n \in N, i$ ranges from $1, 2, \dots, n$.

$$\delta(x_1, x_2, \dots, [x_i, x_i y_i], \dots, x_n) = 0$$

⇒

$$\begin{aligned} & \delta(x_1, x_2, \dots, [x_i, x_i y_i], \dots, x_n) \\ &= \delta(x_1, x_2, x_3, \dots, x_i[x_i, y_i], \dots, x_n) \\ &= \sigma(x_i)\delta(x_1, x_2, x_3, \dots, [x_i, y_i], \dots, x_n) \\ &+ \tau([x_i, y_i])\delta(x_1, x_2, \dots, x_i, \dots, x_n) \\ &= 0 \end{aligned}$$

By the assumption, we have that

$$\tau([x_i, y_i])\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) = 0$$

$$\tau(x_i y_i - y_i x_i)\delta(x_1, x_2, \dots, x_i, \dots, x_n) = 0, \quad \forall x_i, y_i \in N \quad 1 \leq i \leq n$$

As τ is injective, the above equation becomes;

$$\begin{aligned} \tau(x_i y_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \\ &= \tau(y_i x_i)\delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \end{aligned}$$

Then for any $u \in N \exists v \in N$ such that $\tau(u) = v$

$$\begin{aligned} \tau(x_i u)\delta(x_1, x_2, \dots, x_i, \dots, x_n) = \\ \tau(u x_i)\delta(x_1, x_2, \dots, x_i, \dots, x_n) \end{aligned}$$

Now;

$$\begin{aligned}
 0 &= \tau(x_i y_i u - u y_i x_i) \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n). \\
 \forall x_i, y_i, u \in N \quad 1 \leq i \leq n \\
 &= \tau(x_i y_i u - y_i u x_i) \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) \\
 &= \tau(x_i y_i u - y_i x_i u) \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n). \text{ As } \tau \\
 &\text{is an automorphism} \\
 &= \tau(x_i y_i - y_i x_i) \tau(u) \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n), \\
 &\text{since } \tau(u) = v \\
 &= \tau(x_i y_i - y_i x_i) v \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n). \\
 \forall x_i, y_i \quad v \in N \quad 1 \leq i \leq n \\
 \Rightarrow \tau([x_i, y_i]) v \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) &= 0 \\
 \Rightarrow \tau([x_i, y_i]) N \delta(x_1, x_2, x_3, \dots, x_i, \dots, x_n) &= 0
 \end{aligned}$$

By primeness of N and the reality that δ is non-zero left-sided (σ, τ) - n -derivation of N , then $\tau([x_i, y_i]) = 0$. As τ is injective, we have that $[x_i, y_i] = 0$, that is $= 0$

$$\Rightarrow y_i x_i = x_i y_i \forall x_i y_i \in N$$

REFERENCES

- Aroonruviwat, P. and Leerawat, U. (2021). On outer (σ, τ) - n -derivations and commutativity in prime nearrings. *International Journal of Mathematics and Computer Science*, 16(2), 563-575. ijmcs.future-in-tech.net.
- Aroonruviwat, P. and Leerawt, U.(2021). On Commutativity of Nearing with Outer (σ, τ) - n -derivations. *Communications in Mathematics and publications*, 12(1), 161-169, [\[Crossref\]](#)
- Ashraf, M. and Siddeeqe, M. A.(2013). On (σ, τ) - n -derivations in nearrings. *Asian-European Journal of Mathematics*, 6(4),1-14. [\[Crossref\]](#)

Hence N is a commutative ring.

CONCLUSION

In this work, we reviewed some recent papers on (σ, τ) - n -derivation and outer (σ, τ) - n -derivation on left-nearrings, which did not discuss right-nearrings. We considered right-nearrings. The concept of left-sided outer (σ, τ) - n -derivation on right-nearrings is introduced in this study, along with the partial distributive law's proof and the commutativity of nearrings' establishment using the derivation. Further research may be conducted to develop generalized left-sided outer (σ, τ) - n -derivation on right-nearrings and more.

ACKNOWLEDGEMENT

The authors appreciate the assistance from Umaru Musa Yar'adua University Katsina while compiling this work. We are indebted to the reviewers' recommendations, which enable us to improve the entire work thank you all.

- Bell, H.E. and Mason, G. (1987). On derivations in Nearrings and Near-fields, *North-Holland Mathematics studies*, 137, 31–35. [\[Crossref\]](#)
- Clay, J.R. (1992). Nearrings: Geneses and Applications. Illustrated edition, .Oxford Science Publication, USA, pp.3-37. [\[Crossref\]](#)
- Pilz, G. (1983). Nearrings, the theory, and its applications. Second Edition, North-Holland/*American Elsevier*, Mathematics Studies, 23, Amsterdam, pp.6-42. elsevier.com.