

ORIGINAL RESEARCH ARTICLE

New Generalized Odd Fréchet-G (NGOF-G) Family of Distribution with Statistical Properties and Applications.

Ibrahim Abubakar Sadiq¹ , S. I. S. Doguwa¹, Abubakar Yahaya¹ Jamilu Garba¹.¹Department of Statistics, Faculty of Physical Sciences, Ahmadu Bello University, Zaria.**ABSTRACT**

The distribution theory literature contains recent types of parametric distributional models that have been successfully used in the past and whose goodness of fit is sufficient only for certain datasets, suggesting further attention to accommodate a wider range of real-world datasets, for more adaptability, efficiency, and applications. This study aims to develop an extended Fréchet-G family of distributions and study their mathematical properties. The method of Alzaatreh is employed in developing a new lifetime continuous probability distribution called the new Generalized Odd Fréchet-G Family of Distribution. The developed distribution is flexible for studying positive real-life datasets. The statistical properties related to this family are obtained. The parameters of the family were estimated by using a technique of maximum likelihood. A New Generalized Odd Fréchet-Weibull model is introduced. This distribution was fitted with a set of lifetime data. A Monte Carlo simulation is applied to test the consistency of the estimated parameters of this distribution in terms of their bias and mean squared error with a comparison of M.L.E and the maximum product spacing (MPS). The findings of the Monte Carlo simulation show that the M.L.E method is the best technique for estimating the parameter of New Generalized Odd Fréchet-Weibull distribution than the M.PS method. The findings of the application on the data set produce a higher flexibility than some of the competing distributions. In general, our new distributions serve as a viable alternative to other distributions available in the literature for modelling positive data.

ARTICLE HISTORY

Received August 03, 2023.

Accepted September 30, 2023.

Published September 30, 2023.

KEYWORDS

New Odd Fréchet-G Family, Moments, Hazard functions, Maximum Likelihood, Monte Carlo Simulations.

© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

INTRODUCTION

René Fréchet constructed the Fréchet distribution as the maximum value distribution and gave it his name Fisher *et al.* (1928) and Gumbel (1958). As a result of experimental research from numerous disciplines, the Fréchet distribution (FD) is frequently utilized and explored for survival analysis and reliability investigation. The engineering, social, physical, environmental, and life sciences are all fields in which this continuous probability distribution is applicable, Deepshikha *et al.* (2021). In their observations, researchers, such as de Gusmão *et al.* (2011), and Khan *et al.* (2008) noted that the Fréchet distribution is essentially a type II extreme value distribution. The novelty of developing a generalized form of probability distribution drew the attention of academicians and devoted statisticians to the flexibility possessions of the generalized distributions. The generalized extreme value distribution (GEVD) is the result of the Weibull, Gumbel, and Fréchet distribution (GWF) being hybridized. Examples of generalized distribution include the Beta-G, the Gamma-G type I, and the Logistic-G, respectively

developed by Eugene *et al.* (2002), Zografos *et al.* (2005) and Torabi *et al.* (2010). Mc-G developed by Alexander *et al.* (2012), Gamma-G type-3 by Torabi and Montazeri (2012), the Exponentiated-G family by Gupta *et al.* (1998), the Logistic-X family by Tahir *et al.* (2016), the Exponentiated-G class of distribution by Cordeiro *et al.* (2013), Odd Log-logistic by Cordeiro *et al.* (2017) and others like New Weibull-X by Ahmad *et al.* (2018). The T-X family of distributions was first published by Alzaatreh *et al.* (2013), and the exponentiated T-X family of distributions by Alzaghal *et al.* (2013). But Tahir *et al.* (2015) concentrated on the Poisson-X family, whereas Bourguignon *et al.* (2014) presented the Weibull-G family, the Beta Weibull-G by Yousof *et al.* (2017), the Generalized Odd Generalized Exponential-G by Alizadeh *et al.* (2017), the Jamal Weibull-X by Jamal and Nasir (2019), the One Parameter Generalized Odd Fréchet family by Marganpoor *et al.* (2020), the Rayleigh-EOG-X Family by Yahaya and Doguwa (2021), the New Weibull-Odd Fréchet-G Family by Usman *et al.* (2021), the

Correspondence: Ibrahim Abubakar Sadiq. Department of Statistics, Faculty of Physical Sciences, Ahmadu Bello University, Zaria. ✉ isabubakar@abu.edu.ng. Phone Number: +234 813 752 6770.

How to cite: Abubakar Sadiq, I., Doguwa, S. I., Yahaya, A., & Garba, J. (2023). New Generalized Odd Fréchet-G (NGOF-G) Family of Distribution with Statistical Properties and Applications. *UMYU Scientifica*, 2(3), 100 – 107. <https://doi.org/10.56919/usci.2323.016>

Weibull–Odd Fréchet-G Family by Usman *et al.* (2021), the Inverse Lomax-G by Falgore and Doguwa (2020), the One Parameter Odd Fréchet family by Ulhaq and Elgarhy (2018), Gamma-G type-2 by Torabi and Montazeri (2012), Nasir Weibull-G by Jamal and Nasir (2019), Jamal Logistic-X by Jamal and Nasir (2019), etcetera. However, the work of Ul-Haq and Elgarhy (2018), and Marganpoor *et al.* (2020) used one parameter Fréchet family of distribution to describe and model datasets of extreme events. The recent occurrences of extreme events became global issues resulting in hazards to the government administration, lives, and properties. The one-parameter Fréchet family appears sub-optimal and may not perform as its two-parameter version which will provide greater flexibility. Therefore, this work intends to develop and explore the full potential of two parameters Fréchet’s family of distribution for greater flexibility and applicability to extreme events. The two-parameter Fréchet distribution has different parameterizations, and we shall use the one defined by Ahmed *et al.* (2016), a random variable T is said to have a Fréchet distribution with scale parameter α and shape parameter β if its probability density function (pdf) and cumulative distribution function (CDF) are given as,

$$y(t ; \alpha , \beta) = \beta \alpha^\beta t^{-(\beta+1)} \exp \left\{ - \left(\frac{\alpha}{t} \right)^\beta \right\}; 0 < t < \infty, \alpha, \beta > 0 \tag{1}$$

$$Y(t ; \alpha , \beta) = \exp \left\{ - \left(\frac{\alpha}{t} \right)^\beta \right\}; 0 < t < \infty, \alpha, \beta > 0 \tag{2}$$

The Weibull distribution has different parameterizations, and we shall use the one defined by Usman *et al.* (2021). A random variable X is said to have Weibull distribution with scale parameter ϕ and shape parameter ω if its pdf and CDF are given as,

$$f(x ; \phi , \omega) = \phi \omega x^{\omega-1} \exp \{ - (\phi x^\omega) \}; 0 < x < \infty , \phi, \omega > 0 \tag{3}$$

$$F(x ; \phi , \omega) = 1 - \exp \{ - (\phi x^\omega) \}; 0 < x < \infty , \phi, \omega > 0 \tag{4}$$

NEW GENERALIZED ODD FRECHET-G FAMILY OF DISTRIBUTION

We proposed a new family of distribution called the New Generalized Odd Fréchet-G (NGOF-G) family by integrating the density function presented in equation (1) to obtain the CDF given by

$$F_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\} \tag{5}$$

where $\alpha > 0$ is the scale parameter, $\beta > 0$, $\gamma > 0$, is the shape parameter $F_{cdf}(x; \xi)$ is the CDF and ξ the parameters’ vector of the baseline distribution. The corresponding pdf of the CDF in equation (5) is given by

$$f_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \beta \gamma \alpha^\beta f_{pdf}(x; \xi) F_{cdf}^{-(\gamma+1)}(x; \xi) \left[F_{cdf}^{-\gamma}(x; \xi) - 1 \right]^{\beta-1} \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\} \tag{6}$$

where $f_{pdf}(x; \xi)$ is the pdf and ξ is the parameters’ vector of the baseline distribution. Hereafter, a random variable X with density function and distribution function in equations (6) and (5) is denoted by

$$X \sim NGOF - G (\alpha, \beta, \gamma, \xi).$$

Survival and Hazard Rate Function of the NGOF-G Family

The survival function, hazard function, and cumulative hazard function are respectively given as,

$$S_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = 1 - \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\} \tag{7}$$

$$h_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \frac{\beta \gamma \alpha^\beta f_{pdf}(x; \xi) F_{cdf}^{-(\gamma+1)}(x; \xi) \left[F_{cdf}^{-\gamma}(x; \xi) - 1 \right]^{\beta-1} \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\}}{1 - \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\}} \tag{8}$$

$$H_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = - \log \left(1 - \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\} \right) \tag{9}$$

Quantile Function of NGOF-G Family

The quantile function of the NGOF-G family for a simulation through inverting the CDF in equation (5). Supposed that $u[0, 1]$ follows a rectangular distribution, hence, easily derived by the following expressions:

$$x = F_{cdf}^{-\gamma}(x; \xi) \left[\frac{\alpha}{\left(\alpha + (-\log(u))^{\frac{1}{\beta}} \right)} \right] \tag{10}$$

where $F_{cdf}^{-\gamma}(x; \xi)$ is the inverted CDF for every future baseline F and ξ is the parameters’ vector

Suitable Expansion for the PDF and CDF of the NGOF-G Family

Here, we consider the individual terms in the given CDF and pdf of the NGOF-G family presented in equations (5) and (6). Some standard mathematical expansion will be operated upon them, comprising the generalized binomial expansion for negative and positive power, the power series expansion and so on. For instance,

$$\left[e^{-ax} = \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} (ax)^i \text{ and } (1-x)^{ab} = \sum_{j=0}^{\infty} (-1)^j \binom{ab}{j} (x)^j \right],$$

however, equation (5) is the same as,

$$\begin{aligned}
 F_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) &= \exp \left\{ - \left(\alpha (F_{cdf}^{-\gamma}(x; \xi) - 1) \right)^\beta \right\} \\
 &= e^{- \left[\frac{\alpha (1 - F_{cdf}^\gamma(x; \xi))}{F_{cdf}^\gamma(x; \xi)} \right]^\beta} \\
 &= \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} \left[\alpha \left(\frac{1 - F_{cdf}^\gamma(x; \xi)}{F_{cdf}^\gamma(x; \xi)} \right) \right]^{i\beta} \\
 &= \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} \alpha^{i\beta} (1 - F_{cdf}^\gamma(x; \xi))^{i\beta} F_{cdf}^{-\gamma i\beta}(x; \xi) \\
 &= \sum_{i=0}^{\infty} \frac{(-1)^i}{i!} \alpha^{i\beta} \sum_{j=0}^{\infty} \binom{j}{i} (-1)^j (F_{cdf}^{\gamma i\beta}(x; \xi))^j F_{cdf}^{-\gamma i\beta}(x; \xi) \\
 &= \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j}}{i!} \alpha^{i\beta} \binom{j}{i} F_{cdf}^{\gamma i\beta(j-1)}(x; \xi) \tag{11}
 \end{aligned}$$

Therefore, equation (11) reduces to,

$$F_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \sum_{i,j=0}^{\infty} Q_{i,j} F_{cdf}^{\gamma i\beta(j-1)}(x; \xi) \tag{12}$$

where $Q_{i,j} = \frac{(-1)^{i+j}}{i!} \alpha^{i\beta} \binom{j}{i}$

Differentiating equation (12) w.r.t. x we have the corresponding pdf as:

$$f_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \sum_{i,j=0}^{\infty} Q_{i,j} \gamma i\beta (j - 1) f_{pdf}(x; \xi) F_{cdf}^{\gamma i\beta(j-1)-1}(x; \xi) \tag{13}$$

Further simplification of equation (12) is as,

$$F_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \sum_{k=0}^{\infty} b_k A_k(x) \tag{14}$$

where $b_k = \sum_{i,j=0}^{\infty} Q_{i,j}$ and $A_k(x) = F_{cdf}^{\gamma i\beta(j-1)}(x; \xi)$

Differentiate equation (14) w.r.t. x we obtained the corresponding pdf as:

$$f_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) = \sum_{k=0}^{\infty} b_k a_k(x) \tag{15}$$

where $a_k(x) = k f_{pdf}(x; \xi) F_{cdf}^{k-1}(x; \xi)$

Special Sub-Model of the NGOF-G Family

Setting equations (3) and (4) into (5), (6), (7), (8) and (9), then $f(x)$, $F(x)$, $S(x)$, $h(x)$, and $H(x)$ will be for the NGOF-Weibull distribution.

Graph of the Special Sub-Model of the NGOF-G Family

The plot of the density function and the hazard function of the NGOF-Weibull distributions are:





Figure 1: pdf Plot of the New Generalized Odd Fréchet-Weibull Distribution

Figure 1 displays the shapes and the behaviour of the density function of the New Generalized Odd Fréchet-Weibull model at different parameter values. It is also seen how different parameters interact with one another. For instance, in the pdf plots, if the parameters are assumed to equal values, the distribution is symmetrical, whereas it is more positively skewed the greater the difference between the parameters' values and the bell shape, the not greater the difference between the parameters' values.



Figures 2: HF Plot of the New Generalized Odd Fréchet-Weibull Distribution

Figure 2 displays the shapes and the behaviour of the hazard functions of the New Odd Fréchet-Weibull distribution at different parameter values. The graph

shows modified unimodal and modal shapes of hazard rates at different values of the parameters.

Moments of the NGOF-G Family

The role of moments in application to statistics is clear, and the most essential characteristics of a probability model can be examined using moments. Evaluation in statistical inference is necessary, the most vital properties of the distributions were derived using the moments. The rth ordinary moment of a random variable X that follows the New Generalized Odd Frechet-G (NGOF-G) family is given as:

$$\begin{aligned} \mu_r' &= E(X^r) \\ &= \int_0^\infty x^r f_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) dx \\ &= \int_0^\infty x^r \sum_{k=0}^\infty b_k a_k(x) dx \\ &= \sum_{k=0}^\infty b_k \int_0^\infty x^r a_k(x) dx \\ &= \sum_{k=0}^\infty b_k E[Z_k^r] \end{aligned} \tag{16}$$

where $E[Z_k^r] = \int_0^\infty x^r k f_{pdf}(x; \xi) F_{cdf}^{k-1}(x; \xi) dx$

Moment-Generating Function of the NGOF-G Family

The moment-generating function of a random variable X that follows the New Generalized Odd Frechet-G (NGOF-G) family is given as:

$$\begin{aligned} M_X^{NGOF-G}(t) &= E(e^{tx}) \\ &= \int_0^\infty e^{tx} f_{NGOF-G}(x; \alpha, \beta, \gamma, \xi) dx \\ &= \int_0^\infty e^{tx} \sum_{k=0}^\infty b_k a_k(x) dx \\ &= \sum_{k=0}^\infty b_k \int_0^\infty e^{tx} a_k(x) dx \\ &= \sum_{k=0}^\infty b_k E[e^{tZ_k}] \end{aligned} \tag{17}$$

where $E[e^{tZ_k}] = \int_0^\infty e^{tx} k f_{pdf}(x; \xi) F_{cdf}^{k-1}(x; \xi) dx$

Entropies of the NGOF-G Family

The entropy of any random variable X is a measure of indecisiveness, variability, and details properties of the probable results of the variable. The entropy of the NGOF-G family is defined mathematically as:

$$\begin{aligned} I_R(\varpi) &= \frac{1}{1-\varpi} \log \left(\int_0^\infty f_{NGOF-G}^\varpi(x; \alpha, \beta, \gamma, \xi) dx \right) \\ &= \frac{1}{1-\varpi} \log \left(\int_0^\infty (\sum_{k=0}^\infty b_k a_k(x))^\varpi dx \right) \end{aligned} \tag{18}$$

where $\varpi > 0$ and $\varpi \neq 1$

The nth entropy is defined by

$$\begin{aligned} I_{nth}(\varpi) &= \frac{1}{\varpi-1} \log \left(1 - \int_0^\infty f_{NGOF-G}^\varpi(x; \alpha, \beta, \gamma, \xi) dx \right) \\ &= \frac{1}{1-\varpi} \log \left(1 - \int_0^\infty (\sum_{k=0}^\infty b_k a_k(x))^\varpi dx \right) \end{aligned} \tag{19}$$

Order Statistics of the NGOF-G Family

Suppose $X_1, X_2, X_3, \dots, X_n$ is a random sample from the NGOF-G distribution and $X_{i:n}$ represent the ith order statistic, then

$$\begin{aligned} & f_{i:n}(x; \alpha, \beta, \xi) \\ &= \frac{n!}{[(i-1)!(n-i)!]} [f_{NGOF-G}(x; \alpha, \beta, \gamma, \xi)] \\ & \quad \times [F_{NGOF-G}(x; \alpha, \beta, \gamma, \xi)]^{i-1} [1 - F_{NGOF-G}(x; \alpha, \beta, \gamma, \xi)]^{n-i} \\ &= \frac{n!}{[(i-1)!(n-i)!]} \left[\sum_{k=0}^\infty b_k a_k(x) \right] \left[\sum_{k=0}^\infty b_k A_k(x) \right]^{i-1} \left[1 - \sum_{k=0}^\infty b_k A_k(x) \right] \end{aligned} \tag{20}$$

Estimation of Parameters of the NGOF-G Family

Suppose that $x_1, x_2, x_3, \dots, x_n$ are the observed values from the proposed NGOF-G family with parameters α, β, γ . Suppose that $\Phi = [\alpha, \beta, \gamma]^T$ is the $[m \times 1]$ vector of the parameter. The log-likelihood function Φ is expressed by

$$\begin{aligned} \ell_n = \ell_n(\Phi) &= n \log(\beta) + n \log(\gamma) \\ & \quad + n\beta \log(\alpha) \\ & \quad + \sum_{i=1}^n \log[f_{pdf}(x_i; \xi)] \\ & \quad - (\gamma + 1) \sum_{i=1}^n \log[F_{cdf}(x_i; \xi)] \\ & \quad + (\beta - 1) \sum_{i=1}^n \log[F_{cdf}^{-\gamma}(x_i; \xi) - 1] \\ & \quad - \sum_{i=1}^n [\alpha (F_{cdf}^{-\gamma}(x_i; \xi) - 1)]^\beta \end{aligned} \tag{21}$$

Taking the partial derivative of equation (21) w.r.t. the parameters $(\alpha; \beta; \gamma)$ are respectively given as:

$$\frac{\partial \ell_n(\Phi)}{\partial \alpha} = \frac{n\beta}{\alpha} - \sum_{i=1}^n [\alpha (F_{cdf}^{-\gamma}(x_i; \xi) - 1)]^\beta \tag{22}$$

$$\frac{\partial \ell_n(\Phi)}{\partial \beta} = \frac{n}{\beta} + n \ln(\alpha) + \sum_{i=1}^n \ln[F_{cdf}^{-\gamma}(x; \xi) - 1] - \sum_{i=1}^n [\alpha(F_{cdf}^{-\gamma}(x; \xi) - 1)]^\beta \ln[\alpha(F_{cdf}^{-\gamma}(x; \xi) - 1)] \tag{23}$$

$$\frac{\partial \ell_n(\Phi)}{\partial \gamma} = \frac{n}{\gamma} + n \ln(\alpha) + \sum_{i=1}^n \ln[F_{cdf}^{-\gamma}(x; \xi) - 1] + (\beta - 1) \sum_{i=1}^n \frac{f_{pdf}(x; \xi) F_{cdf}^{-(\gamma+1)}(x; \xi)}{(F_{cdf}^{-\gamma}(x; \xi) - 1)} + \sum_{i=1}^n f_{pdf}(x; \xi) F_{cdf}^{-(\gamma+1)}(x; \xi) [\alpha(F_{cdf}^{-\gamma}(x; \xi) - 1)]^\beta \tag{24}$$

The MLEs of the parameters $(\alpha; \beta; \gamma)$, says $(\hat{\alpha}; \hat{\beta}; \hat{\gamma})$ are the simultaneous solution of equations (22), (23), and (24) equating them to zero, i.e. $\frac{\partial \ell_n(\Phi)}{\partial \alpha} = 0; \frac{\partial \ell_n(\Phi)}{\partial \beta} = 0; \frac{\partial \ell_n(\Phi)}{\partial \gamma} = 0$. These equations are

intractable and can only be solved using a numerical iterative method.

RESULT OF THE MONTE CARLO SIMULATION

The vast class of computational algorithms known as "Monte Carlo simulations" uses replicated random sampling to produce numerical results. The basic idea is to employ randomness to address problems that could be theoretically deterministic.

M.L.E and M.P.S Techniques

To evaluate the consistency of the new family's parameters, the simulation study was piloted using the Monte Carlo Simulation technique by computing the bias, variance and mean square error of the estimated parameters from the maximum likelihood estimates and the maximum product spacing estimate. The Simulated data is generated using the quantile function in equation (10) and the likelihood function in equation (21) for different sample sizes $n = 50, 100, 150$ with replicate 200 times each. For the NOGF-Weibull distribution parameter values are $(\alpha, \beta, \gamma, \phi, \omega) = (5.0, 1.0, 2.5, 2.0, 3.0)$.

Table 1: Results of the simulated data from the NGOF-Weibull Distribution.

Sample Sizes	Parameters (Actual Values)	M.L.E. Techniques			M.P.S. Techniques		
		Estimates	Bias	RMSE	Estimates	Bias	RMSE
50	α (5.0)	5.0064	0.0064	0.1763	4.1405	-0.8595	1.0245
	β (1.0)	0.9719	-0.0281	0.1429	0.2224	-0.7776	0.7782
	γ (2.5)	2.6584	0.1584	0.3203	3.5509	1.0509	1.2231
	ϕ (2.0)	1.9829	-0.0171	0.1583	0.5131	-1.4869	1.4908
	ω (3.0)	2.9647	-0.0353	0.2416	4.4028	1.4028	1.4928
100	α (5.0)	5.0006	0.0006	0.0200	4.1942	-0.8058	0.9490
	β (1.0)	0.9992	-0.0008	0.0263	0.2274	-0.7726	0.7730
	γ (2.5)	2.5672	0.0672	0.1722	3.4259	0.9259	1.0724
	ϕ (2.0)	1.9987	-0.0013	0.0427	0.5050	-1.4950	1.4982
	ω (3.0)	3.0004	0.0004	0.0139	4.3642	1.3642	1.4462
150	α (5.0)	5.0000	0.0000	0.0000	4.2464	-0.7536	0.9054
	β (1.0)	1.0000	0.0000	0.0000	0.2285	-0.7715	0.7719
	γ (2.5)	2.5387	0.0387	0.1192	3.3680	0.8680	1.0000
	ϕ (2.0)	2.0000	0.0000	0.0000	0.5002	-1.4998	1.5030
	ω (3.0)	3.0000	0.0000	0.0000	4.3476	1.3476	1.4255

Table 1 presents the results obtained from the Monte Carlo Simulation study. The results indicated that the bias and root mean square error decrease toward zero with an increase in sample size. However, the actual value of the parameters and the estimated values are almost equal at different sample sizes and iterative levels for the M.L.E technique. This proves the consistency of the MLE parameter estimates. For the M.P.S technique, the actual value of the parameters and the estimated values are almost not equal at different sample sizes and iterative levels. This proves the least consistency of the M.P.S parameter estimates. The result also means that the M.L.E technique is the best technique for estimating the

parameter of New Generalized Odd Frechet-Weibull distribution than the M.P.S technique.

APPLICATIONS

Here we used some existing real-life data sets to assess the flexibility of our developed family using Weibull distribution as a baseline.

Some of the Competing Models

Ul-Haq and Elgarhy (2018) introduce a random variable X that is said to follow the Odd Frechet-Weibull distribution if its pdf and CDF are given as

$$f(x; \beta, \phi, \omega) = \beta g(x, \xi) \left((1 - \exp\{-(\phi x^\omega)\}) \right)^{-(\beta+1)} (1 - (1 - \exp\{-(\phi x^\omega)\})^\beta)^{\beta-1} \times \exp\left\{-\left(\frac{1 - (1 - \exp\{-(\phi x^\omega)\})}{(1 - \exp\{-(\phi x^\omega)\})}\right)^\beta\right\}; \quad \forall x, \beta, \phi, \omega > 0 \tag{25}$$

$$F(x; \beta, \phi, \omega) = \exp\left\{-\left(\frac{1 - (1 - \exp\{-(\phi x^\omega)\})}{(1 - \exp\{-(\phi x^\omega)\})}\right)^\beta\right\}; \quad \forall x, \beta, \phi, \omega > 0 \tag{26}$$

Marganpoor *et al.* (2020) introduced a random variable X that is said to follow the Generalized Odd Frechet-Weibull distribution if its pdf and CDF are given as

$$f(x; \alpha, \beta, \phi, \omega) = \alpha \beta (\phi \omega x^{\omega-1} \exp\{-(\phi x^\omega)\}) \left((1 - \exp\{-(\phi x^\omega)\}) \right)^{-\alpha-1}$$

$$((1 - \exp\{-(\phi x^\omega)\})^{-\alpha} - 1)^{\beta-1} \exp\{-(G(x, \phi)^{-\alpha} - 1)^\beta\}; \quad \forall x, \alpha, \beta, \phi, \omega > 0 \tag{27}$$

$$F(x; \alpha, \beta, \phi, \omega) = \exp\{-(G(x, \phi)^{-\alpha} - 1)^\beta\}; \quad \forall x, \alpha, \beta, \phi, \omega > 0 \tag{28}$$

Application on Maximum Annual Flood Discharges Data

The data set was originally reported by Montfort (1970) which represents the Maximum Annual Flood Discharges of North Saskatchewan in units of 1000 cubic feet per second, of the North Saskatchewan River at Edmonton, for 47 years. The data are: 19.885, 20.940, 21.820, 23.700, 24.888, 25.460, 25.760, 26.720, 27.500, 28.100, 28.600, 30.200, 30.380, 31.500, 32.600, 32.680, 34.400, 35.347, 35.700, 38.100, 39.020, 39.200, 40.000, 40.400, 40.400, 42.250, 44.020, 44.730, 44.900, 46.300, 50.330, 51.442, 57.220, 58.700, 58.800, 61.200, 61.740, 65.440, 65.597, 66.000, 74.100, 75.800, 84.100, 106.600, 109.700, 121.970, 121.970, 185.560.

Table 2: Parameters Estimates and Goodness of Fit Measures for Maximum Annual Flood Discharges Data

Model	Parameter Estimates and Goodness of Fit					LL	AIC
	$\hat{\alpha}$	$\hat{\beta}$	$\hat{\gamma}$	$\hat{\phi}$	$\hat{\omega}$		
NGOFWD	0.3144	2.1499	0.6957	0.0065	0.6619	179.2229	368.4458
GOFWD	2.0973	4.9876	0.5524	0.2328	-	215.0636	438.1276
OFWD	1.2325	8.3788	1.4295	-	-	1918.734	3831.468
EtWD	2.3463	1.8221	7.2015	-	-	266.6662	489.6832
WD	0.0007	1.7724	-	-	-	225.7065	455.4135
FD	35.246	2.4462	-	-	-	215.1136	434.2272

Table 2 provides the parameter estimates and goodness of fit measures for the New Generalized Odd Frechet-Weibull distribution with other competing models using the maximum annual flood discharges dataset. Akaike's Information Criterion (AIC), Bayesian Information Criterion (BIC) and Hannan-Quinn Information Criterion (HQIC) are the performance metrics. A distribution with the lowest information or performance metrics is regarded as the best in terms of goodness of fit. The new generalized odd Frechet-Weibull distribution is the best model that outperforms other competitors based on the data set.




Figure 3: Histogram Plots of the Distribution of Maximum Annual Flood Discharges Data

CONCLUSION

In this paper, we developed the New Generalized Odd Frechet-G (NGOF-G) family of distributions. The statistical properties of this family comprising the survival function, hazard function, cumulative hazard function, moments, moment generating function, entropies, order

statistics, and MLE are obtained. We further plot the pdf and the hazard rate function to assess the shapes and behaviour of the models at different parameter values. Simulation studies were carried out for the assessment of the consistents of MLE and MPS of the parameters. We present the application of the NGOF-W distribution to the data representing Maximum Annual Flood Discharges employing Weibull as the baseline distribution. The analysis indicated that the NGOFW is the “best fit” model for the Maximum Annual Flood discharge data.

REFERENCES

- Ahmad, Z., Ampadu, C. B., Hamedani, G. G., Jamal, F., & Nasir, M. A. (2019). The New Exponentiated TX Class of Distributions: Properties, Characterizations and Application. *Pakistan Journal of Statistics and Operation Research*, 941-962. [[Crossref](#)]
- Ahmad, Z., Elgarhy, M., & Hamedani, G. G. (2018). A New Weibull-X Family of Distributions: Properties, Characterizations and Applications. *Journal of Statistical Distributions and Applications*, 5(1), 1-18. [[Crossref](#)]
- Ahmed, Haitham, Cordeiro, Edwin and Zohdy (2016). “The Weibull Fréchet Distribution and Its Applications”. *Journal of Applied Statistics*, [[Crossref](#)]
- Alexander, C., Cordeiro, G. M., Ortega, E. M., & Sarabia, J. M. (2012). Generalized Beta-Generated Distributions. *Computational Statistics & Data Analysis*, 56(6), 1880-1897. [[Crossref](#)]
- Alizadeh, M., Ghosh, I., Yousof, H. M., Rasekhi, M., & Hamedani, G. G. (2017). The Generalized Odd Generalized Exponential Family of Distributions: Properties, Characterizations and Application. *Journal of Data Science*, 15(3), 443-465. [[Crossref](#)]
- Alzaatreh, A., Lee, C., Famoye, F. (2013b). A New Method for Generating Families of Continuous Distributions. *Metron*, 71, 63–79. [[Crossref](#)]
- Alzaghal, A., Famoye, F. and Lee, C. (2013): Exponentiated T-X Family of Distributions with some Applications. *International Journal of Statistics and Probability*, 2(3), 31–49. [[Crossref](#)]
- Bourguignon, M., Silva, R. B. and Cordeiro, G. M. (2014). The Weibull-G Family of a Probability Distribution. *Journal of Data Science*, 12, 53-68. [[Crossref](#)]
- Chukwu A. U., & Ogunde A. A., (2015). On the Beta Mekaham Distribution and Its Application. *American Journal of Mathematics and Statistics* 2015, 5(3): 137-143. [[Crossref](#)]
- Cordeiro, G. M., Alizadeh, M., Ozel, G., Hosseini, B., Ortega, E. M. M., & Altun, E. (2017). The Generalized Odd Log-Logistic Family of Distributions: Properties, Regression Models and Applications. *Journal of Statistical Computation and Simulation*, 87(5), 908-932. [[Crossref](#)]
- Cordeiro, G. M., Ortega, E. M., & da Cunha, D. C. (2013). The Exponentiated Generalized Class of Distributions. *Journal of data science*, 11(1),127. [[Crossref](#)], [[Crossref](#)]
- De Gusmao, F. R., Ortega, E. M., & Cordeiro, G. M. (2011). The generalized inverse Weibull distribution *Statistical Papers*, 52, 591-619. [[Crossref](#)]
- Deepshikha, D.; Bhatia, D.; Bhupen, k. B.; Bhupen, B. (2021). “Some Properties on Fréchet-Weibull Distribution with Application to Real Life Data. *Mathematics and Statistics*. 9(1): 8-15. [[Crossref](#)]
- Eugene, N., Lee, C. and Famoye, F. (2002). Beta-Normal Distribution and It Applications. *Communications in Statistics: Theory and Methods*, 31: 497-512. [[Crossref](#)]
- Falgore, J. Y., & Doguwa, S. I. (2020). The Inverse Lomax-g Family with Application to Breaking Strength Data. *Asian Journal of Probability and Statistics*, 49-60. [[Crossref](#)]
- Fisher, R. A.; Tippett, L. H. C. (1928). "Limiting Forms of the Frequency Distribution of the Largest and Smallest Member of a Sample". *Proceedings of the Cambridge Philosophical Society*. 24 (2): 180–190. [[Crossref](#)]
- Gumbel, E. J. (1958). *Statistics of Extremes*. New York: *Columbia University Press*. OCLC 180577. [[Crossref](#)]
- Gupta, R. C., Gupta, P. L. and Gupta, R. D. (1998). Modelling Failure Time Data by Lehmann Alternatives. *Communications in Statistics - Theory and Methods* 27, 887-90. [[Crossref](#)]
- Haight, Frank A. (1967), *Handbook of the Poisson Distribution*, New York, NY, USA: *John Wiley & Sons*, ISBN 978-0-471-33932-8.

- Jamal, F., Ahmad, Z., Ampadu, C. B., Hamedani, G. G., & Nasir, M. A. (2019). The New Exponentiated TX Class of Distributions: Properties, Characterizations, and Application, *Pakistan Journal of Statistics and Operation Research*, 941–962. [[Crossref](#)]
- Jamal, F., Arslan Nasir, M., Ozel, G., Elgarhy, M., & Mamode Khan, N. (2019). Generalized Inverted Kumaraswamy Generated Family of Distributions: Theory and Applications, *Journal of Applied Statistics*, 46(16), 2927–2944. [[Crossref](#)]
- Khan M.S.; Pasha G.R.; Pasha A.H. (February 2008). "Theoretical Analysis of Inverse Weibull Distribution". *WSEAS Transactions on Mathematics*. 7 (2). pp. 30–38.
- Marganpoor, S., Ranjbar, V., Alizadeh, M., & Abdollahnezhad, K. (2020). Generalised Odd Frechet Family of Distributions: Properties and Applications. *Statistics in Transition new series*, 21(3), 109-128. [[Crossref](#)]
- Montfort, M. A. On Testing that the Distribution of Extremes is of Type I when Type II is the Alternative, 11, 421–427, 1970. [[Crossref](#)]
- Oguntunde P. E., A. O. Adejumo & K. A. Adepoju (2016). Assessing the Flexibility of the Exponentiated Generalized Exponential Distribution, *Pacific Journal of Science and Technology*, 17(1), 49-57
- Tahir, M. H., Cordeiro, G. M., Alizadeh, M., Mansoor, M., Zubair, M., & Hamedani, G. G. (2015). The Odd Generalized Exponential Family of Distributions with Applications. *Journal of Statistical Distributions and Applications*, 2(1), 1-28. [[Crossref](#)]
- Tahir, M. H., Cordeiro, G. M., Alzaatreh, A., Mansoor, M., & Zubair, M. (2016). The logistic-X Family of Distributions and Its Applications. *Communications in Statistics-Theory and Methods*, 45(24), 7326-7349. [[Crossref](#)]
- Tahir, M. H., Zubair, M., Cordeiro, G. M., Alzaatreh, A. and Mansoor, M. (2016). The Poisson-X Family of Distributions. *Journal of Statistical Computation and Simulation*, 86, 2901- 2921. [[Crossref](#)]
- Tahir, M. H., Zubair, M., Mansoor, M., Cordeiro, G. M., ALIZADEHK, M., & Hamedani, G. (2016). A New Weibull-G Family of Distributions. *Hacettepe Journal of Mathematics and Statistics*, 45(2), 629-647. [[Crossref](#)]
- Thomas W. Keelin (2016) The Meta log Distributions. *Decision Analysis* 13(4):243-277. <http://dx.doi.org/10.1287/deca.2016.0338>. [[Crossref](#)]
- Torabi, H. and Montazari, N.H. (2012). The gamma-uniform distribution and its application. *Kybernetika* 48:16–30.
- Torabi, H., & Bagheri, F. L. (2010). Estimation of Parameters for an Extended Generalized Half Logistic Distribution Based on Complete and Censored Data
- Ul Haq, M. A., & Elgarhy, M. (2018). The odd Frèchet-G Family of Probability Distributions. *Journal of Statistics Applications & Probability*, 7(1), 189-203. [[Crossref](#)]
- Usman A., Doguwa S. I. S., Alhaji B. B., and Imam A. T. A New Weibull- Odd Frechet G Family of Distributions. *Transactions of the Nigerian Association of Mathematical Physics* Volume 14, (January - March 2021 Issue), pp133 –142
- Yahaya, A., & Doguwa, S. I. S. Theoretical Study of Rayleigh-Exponentiated Odd Generalized-X Family of Distributions. *Transactions of the Nigerian Association of Mathematical Physics* Volume 14, (January-March, 2021 Issue), pp143–154© *Trans. of NAMP on*
- Yousof, H. M., Rasekhi, M., Afify, A. Z., Ghosh, I., Alizadeh, M., & Hamedani, G. G. (2017). The Beta Weibull-G Family of Distributions: Theory, Characterizations and Applications. *Pakistan Journal of Statistics*, 33(2). [[Crossref](#)]
- Zografos, K., & Balakrishnan, N. (2009). On Families of Beta-Generalized Gamma-Generated Distributions and Associated Inference, *Statistical Methodology*, 6(4), 344–362. [[Crossref](#)]