

REVIEW ARTICLE

The Potentials of Biosurfactants as Anti-Inflammatory and Anti-Viral Agents Against Covid-19: A Mini Review

Lawal, Ibrahim*¹, Fardami, Aminu.Yusuf², Bello, Suleiman³, Aisha, Habibu⁴ and Zainab, Muhammad Sunusi²

¹Microbiology Department, College of Natural and Applied Sciences, Al-Qalam University Katsina, Nigeria

²Department of Microbiology, Faculty of Chemical and Life Science, Usmanu Danfodio University Sokoto, Nigeria

³Department of Bioscience, Faculty of Science, Universiti Teknologi Malaysia, Johor Bahru, Johor, Malaysia.

⁴Department of Science Laboratory Technology, College of Science and Technology, Umaru Ali Shinkafi Polytechnic Sokoto, Nigeria

ABSTRACT

Infection due to strain of severe acute respiratory syndrome coronavirus 2 (SARS COV2) has grown to be of global public health significance. Biotechnology uses living organisms such as microbes to produce metabolites like biosurfactants. Biosurfactants are amphiphilic surface active biomolecules that were proven to have therapeutic function against some groups of microbes including viruses. They also have anti-inflammatory potential through their interaction with viral membranes and macromolecules to decrease cytosolic phospholipase A2, which is the beginning of an anti-inflammatory response, and are recognized structurally by toll-like receptors (TLR-2), which are released when neutrophils are stimulated. They can also play vital role in aiding the human body to have inflammatory response. The functional groups of biosurfactants interact with the viruses membrane structure. Some groups of biosurfactants cause physicochemical processes that render viruses inactive. Therefore it can generally be understood that biosurfactants destroy the virus's envelope and the viral membrane's structures. The principle behind biosurfactant's anti viral property is due to the hydrophilic properties that are within the acetyl groups. Additionally, the hydrophobic properties of biosurfactant are also important in making it to have antiviral activity. These activities of biosurfactants against viruses make it to be potential anti-inflammatory and anti-viral agents against Covid-19. Therefore this paper is aimed to produce a mini review on the anti-inflammatory and anti-viral potential against Covid-19. And the review also highlights some of the desirable properties and benefits of biosurfactants as anti-corona viruses.

ARTICLE HISTORY

Received November 19, 2022

Accepted December 16, 2022

Published December 30, 2022

KEYWORDS

Biosurfactants, Anti-inflammatory, Anti-viral, COVID-19, Interleukins

© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

INTRODUCTION

The surface and interfacial tension between different media are reduced by surfactants, which are surface-active compounds with dual characteristics having both water-loving and water-hating functional groups. These agents are classified into chemical and biological surfactants based on their origin (Santos *et al.*, 2017). They are also classified into glycolipids, phospholipids, lipopeptides, as well as higher molecular weight or polymeric surfactants based on monomers they are composed of (Fardami *et al.*, 2022). Biosurfactants are secondary metabolites produced by a variety of microorganisms, including bacteria, molds and yeasts. They are categorized according to their structural components and the producing microbes such as *Pseudomonas aeruginosa*, *Bacillus subtilis*, and *Lactobacillus* sp., *Candida bombicola*, *Torulopsis bombicola*, *Penicillium*

chrysogenum and *Aspergillus versicolor* (Perfumo *et al.*, 2017; Fardami *et al.*, 2022). During growth, they are either produced within the cells or excreted in medium (Santos *et al.*, 2016).

Biosurfactants are a diverse group of surface-active compounds produced by microorganisms (Inès *et al.*, 2023). These compounds have the ability to reduce surface tension and increase the solubility of hydrophobic compounds in aqueous solutions (Patel *et al.*, 2023). Biosurfactants have a wide range of applications, including in the oil and gas industry, environmental remediation, and bioremediation (Singh, 2023). Recently, there has been growing interest in the use of biosurfactants as antiviral agents, particularly in the context of inhibiting viral infection (Kisla *et al.*, 2023).

Correspondence: Lawal, Ibrahim. Microbiology Department, College of Natural and Applied Sciences, Al-Qalam University Katsina-Nigeria. ✉ ibrahimlawal.mcb@auk.edu.ng Phone Number; +2348034464896

How to cite: Lawal, Ibrahim, Fardami, Aminu.Yusuf, Bello, Suleiman, Aisha, Habibu and Zainab, Muhammad Sunusi (2023). The Potentials of Biosurfactants as Anti-Inflammatory and Anti-Viral Agents Against Covid-19: A Mini Review. UMYU Scientifica, 1(2), 188 – 194. <https://doi.org/10.56919/usci.1222.019>

Viral infections remain a significant health concern worldwide, and traditional antiviral therapies can have limitations such as toxicity, resistance, and high cost (Sokhela *et al.*, 2023). Biosurfactants offer a potential alternative approach to inhibiting viral infections (Ukaegbu *et al.*, 2023). Studies have shown that certain biosurfactants can disrupt the lipid envelope of enveloped viruses, leading to viral inactivation or inhibition of viral replication. The lipid envelope is a critical component of many viruses, and disruption of the envelope can prevent the virus from entering host cells or replicating (Tripathy *et al.*, 2023).

Biosurfactants have been shown to be effective against a range of viruses, including human immunodeficiency virus (HIV), herpes simplex virus (HSV), influenza virus, and respiratory syncytial virus (RSV) (El Khalloufi and Oudra, 2023; Sil *et al.*, 2023; Antony *et al.*, 2022). For example, rhamnolipids, a type of biosurfactant produced by *Pseudomonas aeruginosa*, have been shown to inhibit the entry and replication of HIV in human cells (Vanreppelen *et al.*, 2023). Similarly, sophorolipids, another type of biosurfactant produced by yeast, have been shown to inhibit the entry and replication of HSV in human cells (Karnwal *et al.*, 2023).

In December 2019, a new strain of coronavirus that causes the coronavirus illness 2019 (COVID-19) developed and spread globally. A global public health emergency has emerged because of the COVID-19 in different countries of the world (Venugopal *et al.*, 2020). Apart from severe acute respiratory syndrome coronavirus 2 (SARS-CoV-1) the brand-new strain of the coronavirus with rapid human-to-human transmission, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused a global public health disaster and claimed many lives (Chen *et al.*, 2019; Chan *et al.*, 2020). After contracting the infection, the infected individuals primarily depend on their immunity to fight the virus, with supportive therapy being recommended if difficulties arise (Grasselli *et al.*, 2020). According to research, the expression of angiotensin-converting enzyme 2 (ACE2) in host cells is the initial stage in disease initiation process of the new coronavirus strain (Hoffmann *et al.*, 2020).

The defense system sends a lot of cells to kill the virus once it gets inside the lungs (Huang *et al.*, 2020). The system becomes overactive once the cytokine storm (an immune reaction in which many cytokines are released into the blood too quickly) forms and kills the healthy cells (Mahalaxmi *et al.*, 2020). In addition to the virus' ability to defy therapy, the immune system presents a significant challenge when choosing the best course of action and vaccine. The immune system's ability to maintain balance is weakened by SARS-CoV-2, which also causes the cytokine storm. The cytokine storm has been associated with the considerable challenges seen in COVID-19

patients. After attaching, the spike protein splits in two, causing a conformational shift that makes it easier for the virus to fuse and enter the cell. The stages of viral replication in cells were recently thoroughly described by Naughton *et al.* (2019). Over the past ten years, the number of BSs with a medical focus has increased. Biosurfactant fights a variety of diseases because of its antiviral, antibacterial, and antifungal properties (Naughton *et al.*, 2019). The use of biosurfactants as antiviral agents is still in the early stages of development, and more research is needed to fully understand their potential applications. This mini review is aimed to highlight the potential application of biosurfactants as anti viral and anti inflammatory agents.

ANTI-INFLAMMATORY POTENTIAL OF BIOSURFACTANTS

Phospholipase A2 (PLA2) facilitates the secretion of arachidonic acid (AA). Different PLA2 variants are collectively referred to as cytosolic phospholipase-A2 (cPLA2). An inflammatory response occurs as arachidonic acid is released and changed into inflammatory mediators. Eicosanoids, which are generated as a result of arachidonic acid, are first created in order to maintain the inflammatory process. Biosurfactants interact with cell membranes and macromolecules to decrease cytosolic phospholipase A2, which is the beginning of an anti-inflammatory response, and are recognized structurally by toll-like receptors (TLR-2), which happened when neutrophils in an in vitro model were stimulated by *Rhodococcus ruber's* trehalolipids to release inflammatory cytokines (Baeva *et al.*, 2014). The lipids were being supplied by these neutrophils. According to Chereshevnev *et al.* (2010) and Gein *et al.* (2011), induced *R. ruber* glycolipids in mononuclear cells facilitate the synthesis of IL-12, IL-18, and reactive oxygen species (ROS) and increase the production of TNF- α , IL-1b, and IL-6. In rat and fish models, surfactin administration decreased levels of pro-inflammatory cytokines while raising levels of anti-inflammatory cytokines (Ramasami *et al.*, 2009; Giri *et al.*, 2016). According to research by Zhang *et al.* (2015), the antibacterial lipopeptide biosurfactant surfactin was found to inhibit lipopolysaccharide-induced signaling pathways, impair macrophage function, prevent IL-12 expression, and reduce the expression of the toll-like receptor-4 protein, which increased the anti-inflammatory effect.

According Park *et al.* (2013), *Staphylococcus aureus's* surfactin was able to considerably decrease the proinflammatory mediators which also raised STAT-3 phosphorylation, preventing the synthesis and release of haeme oxygenase-1 and blocked the lipoteichoic acid-induced signaling pathway (HO-1). The impact of biosurfactants produced by yeast species with anti-inflammatory activity was also

indicated by a few small investigations that were carried out. Immunoglobulin E (IgE) levels, IL-6, STAT3, and TLR-2 mRNA expression, as well as lung inflammation, were all reduced by sophorolipids (SLs) from *Candida bombicola* (Hagler *et al.*, 2007; Vakil *et al.*, 2013). Therefore, the study showed that sophorolipids downregulate the IgE coding genes, acting as an anti-inflammatory molecule and possible therapeutic substance (Hagler *et al.*, 2007; Bluth *et al.*, 2006).

In an in-vivo experiment, sophorolipids were found to have anti-inflammatory properties and reduced sepsis-related mortality in a rat model (Hardin *et al.*, 2007; Mueller *et al.*, 2006). Similar to this, SLs reduced nitric oxide (a reactive oxygen species), regulated inflammatory responses, and enhanced survival rates in a different rat model research (Bluth *et al.*, 2006). It has been shown that both natural and synthesized sophorolipids have strong anti-inflammatory, spermicidal, and anti-HIV activity (Shah *et al.*, 2005). The SLs inhibit the expression of inflammatory cytokines (Mueller *et al.*, 2006), and these results suggest that SLs would be a promising therapeutic agents for immunomodulation or anti-inflammatory in disorders that are chronically inflammatory. *Pseudomonas antarctica* secretes mannosylerythritol lipids, which have anti-inflammatory effects since they have suppressed the inflammatory mediators (Morita *et al.*, 2011).

MECHANISM OF TISSUE DAMAGE AND INFLAMMATORY RESPONSE OF THE BODY AGAINST COVID 19

The body's defense system promptly mobilizes a huge quantities of cells to fight the virus, especially by securing the help of lymphocytes, after the virus has entered the vertebrate host cell via (Yang, 2020). High levels of cytokine storm have been found to be present in COVID-19-positive individuals, and these levels are also related to the viral load in these patients (Wang *et al.*, 2020). When a cytokine storm develops, the immune system becomes overactive and kills healthy cells (Mahalaxmi *et al.*, 2020). Furthermore, larger lymphocyte and IL-6 levels unintentionally cause more pulmonary injury (Akhmerov and Marban, 2020). Furthermore, while contemplating the best course of therapy and vaccination possibilities, the virus's capacity to elude the immune system poses serious challenges. The resultant harm may be brought on by direct SARS-CoV-2 infection of cells, hypoxia brought on by pulmonary injury brought on by immunological reactions (Harshada, 2014).

In the course of the cytokine storm, excessive levels of cytokines like IL-1b and IL-18 are created, which may permanently harm different organs. The biosurfactants are well known for their significant contribution to the body's defense against disease causing microorganisms and their ability to reduce inflammation (Sajid *et al.*, 2020). Different anti-microbial disorders have been successfully treated with the help of the glycolipid and lipopeptide

types of biosurfactants (Liu and Li, 2020). One of its subtypes, surfactin, a naturally occurring cyclic lipopeptide, has been demonstrated to possess a variety of microbial activities (Singh and Cameotra, 2004). These activities are started by blocking the signaling of cell survival, platelet aggregation, and lowering the cytokine storm by putting forth anti-inflammatory effects.

In order to reduce the effects of the cytokine storm brought on by viral infection in the infected individuals, it may be possible to employ surfactants of biological origin. Speculative mode of action for biological surfactant's ability to reduce inflammation in the COVID-19 illness has been speculated. The Spike of SARS-CoV-2 is split in two upon contact, which causes a conformational shift that facilitates the attachment of the virus and its penetration into the cell. Viral proteins N, S, 3a, and 7a stimulate nuclear factor Kappa beta pathway, a common pathway involved in numerous diseases. Pro-IL-1b and procaspase-1 are both transcriptionally regulated by NF- κ B, which, upon activation, moves into the nucleus. Pro-IL-1b and procaspase 1 are split into IL-1b and caspase when other signals, such as elevated Ca²⁺ and reactive oxygen species, are recognized. This triggers the release of many mediators, which culminates in a cytokine storm causing cell destruction and death. Since heme is necessary for the creation of biliverdin, ferrous iron, and carbon monoxide, which might reduce inflammation and stress brought on by SARS-CoV-2 viral infection, it has been discovered that COVID-19 patients have heme production that is inhibited (Fujioka *et al.*, 2017; Takeda *et al.*, 2017; Saimmai *et al.*, 2020). By activating the HO-1 and TH1 macrophage cells, the biological surfactant may reduce the production of NF- κ B if given to COVID-19 patients (Rodrigues *et al.*, 2006).

This would therefore lessen the production of cytokines including TNF α , IL-1B, IL-6, and IL-2, which would lessen the impact of the cytokine storm in COVID-19 patients. Even though biosurfactants are naturally formed, they are renowned for their emulsifying abilities in pharmaceuticals and vaccines. Therefore, when combined with traditional antigens for the treatment of COVID-19 disease, these surfactants contain immunological adjuvants that are both non-toxic and non-pyrogenic (Paulino *et al.*, 2016). Consequently, these demonstrate that surfactants of biological origin play a significant role as immunosuppressive drugs and may be utilized extensively as a combine therapy toward reducing the inflammatory reactions brought on by coronavirus infection.

BIOSURFACTANTS' ANTI-VIRAL ACTIVITY AGAINST COVID-19

A few biosurfactants cause physiochemical processes that render viruses inactive (Vollenbroich *et al.*, 1997). Generally, it is claimed that biosurfactants destroy the virus's envelope and the viral membrane's structures (Shah

et al., 2005). The biological surfactant's hydrophilic properties are caused by the acetyl groups, which encourage antiviral action (Borsanyiova *et al.*, 2016). Additionally, the hydrophobic properties with a particular amount of carbon atoms neutralize the virucidal actions (Kracht *et al.*, 1999). High viral inactivation was experienced for surfactants having fatty acid chain longer than fifteen carbon atom; same was reported for monomethyl esters in the semliki forest virus (Kracht *et al.*, 1999). The activity of biological surfactants against coronavirus has been confirmed, and patents on several viruses have been secured (Bonvial *et al.*, 2009; Gross *et al.*, 2014; Gross *et al.*, 2004; Borzeix, 1999; Gross and Shah, 2007). Since SARS-CoV-2 is an enveloped virus, it is possible to apply the evidence from these experiments to it, and the following mechanism of action has been described.

The functional groups of biosurfactants interact with the cell membrane of the virus as the virus enters the host cell and move into bilayered lipid membrane, changing the permeability by either forming ion channels or disrupting the membrane system (Subramaniam *et al.*, 2023). High concentrations of BSs cause the capsid and its surrounding envelope to completely disintegrate (Sarangi *et al.*, 2023). The spike protein and lipid envelope disturbances are enclosed into micelles, which renders the virus inactive (Nitschke and Marangon, 2022). The created micelle has the capacity to serve as carrier that could transport the medication to the site of infection while also providing protection under risky circumstances (Nakashini *et al.*, 2009). Therefore, the ability of biosurfactants to form micelles would make them efficient drug delivery agents for the treatment of covid-19 infection. Additionally, it inactivates the effects of the virus before adsorption or penetration but does not alter viral reproduction.

DESIRABLE PROPERTIES AND BENEFITS OF BIOSURFACTANTS AS ANTI-CORONA VIRUSES

As a more effective means of taking preventative measures or receiving treatment for the SARS-CoV-2 infection,

REFERENCES

- Akhmerov, A. and Marbán, E. (2020). COVID-19 and the heart. *Circ Research*, 126:1443-1455. [Crossref]
- Antony, S., Sukumaran, T. U., Rathinam, P., Reshmy, R., Binod, P., Pandey, A., & Sindhu, R. (2022). *Biosurfactants in respiratory viruses and the Coronavirus disease 2019 pandemic*. In Green Sustainable Process for Chemical and Environmental Engineering and Science (pp. 439-450). [Crossref] Academic Press.
- Baeva, T., Gein, S., Kuyukina, M., Ivshina, I., Kochina, O and Chereshevnev. V. (2014). Effect of glycolipid

recommendations of some benefits that will be derived when biosurfactants are used will be made in this paper.

1. Surfactants of biological origin should be used for various of purposes in industries including food, medicine, cosmetics, detergents, pharmaceuticals and others. However, its ability to reduce inflammation might be a novel approach to various ways of treating infections due to coronavirus
2. Hand hygiene could serve as shield from virus. A more effective defence in preventing viral infection is the use of biosurfactants-encoded handwash or hand sanitisers.
3. The amphiphilic properties of the biosurfactant make it simpler to interface with the lipid bilayer of SARS-CoV-2 and would allow for the easy clearance of the virus by allowing the destruction of the viral DNA.
4. Because of their strong propensity for medication distribution, BSs are particularly well-suited for the job. Because they have a protective antiviral and anti-inflammatory potential against the SARS-CoV-2 virus, it is highly advised to utilize or produce any drug from BSs in addition to conventional medications or vaccinations.
5. Incorporating some therapeutic plants (i.e when using herbal preparations) with microbial surfactants has great promise for effectively removing the viral burden from the human body.

CONCLUSION

The use of biosurfactant as anti-inflammatory in COVID-19 patients, as well as anti-viral agent in the inhibition of viral replication can serve as a potential source of treating Covid 19 where natural compounds produced by microorganisms have shown to have anti-inflammatory and anti-viral properties. The potential of biosurfactants as anti-inflammatory and anti-viral agents against COVID-19, and their potential applications in the management of this disease is promising.

Rhodococcus biosurfactant on secretory activity of neutrophils in vitro. *Bulletin of Experimental Biology and Medicine*, 157:238. [Crossref]

- Bluth, M., Smith-Norowitz, T., Hagler, M., Beckford, R., Chice, S. and Shah, V. (2006). Sphorolipids decrease IgE production in U266 cells. *Journal of Allergy and Clinical Immunology*, 117. S202. [Crossref]

- Bluth, M.H., Kandil, E., Mueller, C.M., Shah, V., Lin, Y-Y. and Zhang, H. (2006). Sphorolipids block lethal effects of septic shock in rats in a cecal ligation and puncture model of experimental sepsis. *Critical Care and Medicine*, 34. [Crossref]

- Bonvila, X.R., Roca, S.F. and Pons, R.S.(2009). Inventors; NOVACYT, assignee. Antiviral use of cationic surfactant. *United States patent application*, 12/375:774.
- Borsanyiova, M., Patil, A., Mukherji, R., Prabhune, A. and Bopegamage, S. (2016). Biological activity of sophorolipids and their possible use as antiviral agents. *Folia Microbiology*, 61:85-89. [[Crossref](#)]
- Borzeix, C.F. (1999). Use of sophorolipids comprising diacetyl lactones as agent for stimulating skin fibroblast metabolism. *Patent WO99/62479*.
- Chan, J.F-W., Yuan, S., Kok, K-H., To, K-K-W., Chu, H. and Yang, J. (2020). A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet*, 395:514-523. [[Crossref](#)]
- Chen, N., Zhou, M., Dong, X., Qu, J., Gong, F. and Han, Y. (2020). Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*, 395:507-513. [[Crossref](#)]
- Chereshnev, V., Gein, S., Baeva, T., Galkina, T., Kuyukina, M. and Ivshina, I. (2010). Modulation of cytokine secretion and oxidative metabolism of innate immune effectors by *Rhodococcus* biosurfactant. *Bulletin of Experimental Biology and Medicine*, 149:734. [[Crossref](#)]
- El Khalloufi, F., & Oudra, B. (2023). Seaweeds as a New Source of Bioactive Compounds and Potent Biotechnological Applications. In *Seaweed Biotechnology* (pp. 229-287). Apple Academic Press. [[Crossref](#)]
- Fardami, A.Y., Kawo, A.H., Yahaya, S., Lawal, I., Sani, A.A. and Maiyadi, K.A. (2022). A Review on Biosurfactant Properties, Production and Producing Microorganisms. *Journal of Biochemistry, Microbiology and Biotechnology*, 10(1): 5-12. [[Crossref](#)]
<https://doi.org/10.54987/jobimb.v10i1.656>
- Fujioka, K., Kalish, F., Zhao, H., Lu, S., Wong, S. and Wong, R.J.(2017). Induction of heme oxygenase-1 attenuates the severity of sepsis in a non-surgical preterm mouse model. shock: injury, inflammation, and sepsis. *Laboratory Clinical Approaches*, 47:242-250. [[Crossref](#)]
- Gein, S., Kuyukina, M., Ivshina, I., Baeva, T., Chereshnev, V. (2011) In vitro cytokine stimulation assay for glycolipid biosurfactant from *Rhodococcus ruber*: role of monocyte adhesion. *Cytotechnology*, 63:559-566. [[Crossref](#)]
- Giri, S.S., Sen, S.S., Jun, J.W., Sukumaran, V., Park, S.C.(2016) Role of Bacillus subtilis VSG4-derived biosurfactant in mediating immune responses in *Labeorohita*. *Fish Shell fish Immunology*, 54: 220-229. [[Crossref](#)]
- Grasselli, G., Zangrillo, A., Zanella, A., Antonelli, M., Cabrini, L. and Castelli, A.(2020). Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. *Journal of American Medical Association*, 323: 1574-1581. [[Crossref](#)]
- Gross, R.A. and Shah, V. (2007). Anti-herpes virus properties of various forms of sophorolipids. Patent US. 200031243: A2
- Gross, R.A., Shah, V. and Doncel, G. (2004). Spermicidal and virucidal properties of various forms of sophorolipids. Patent US. 20040242501:A1.
- Gross, R.A., Shah, V. and Doncel, G. (2014). Virucidal properties of various forms of sophorolipids. Patent US, 8648055:B2.
- Hagler, M., Smith-Norowitz, T., Chice, S., Wallner, S., Viterbo, D. and Mueller, C.(2007). Sophorolipids decrease IgE production in U266 cells by downregulation of BSAP (Pax5), TLR-2, STAT3 and IL-6. *Journal of Allergy and Clinical Immunology*, 119. 263. [[Crossref](#)]
- Hardin, R., Pierre, J., Schulze, R., Mueller, C.M., Fu, S.L. and Wallner S.R. (2007). Sophorolipids improve sepsis survival: effects of dosing and derivatives. *Journal of Surgical Research*, 142:314-319. [[Crossref](#)]
- Harshada, K. (2014). Biosurfactant: a potent antimicrobial agent. *Journal of Microbiology Experiment*, 1:173-177. [[Crossref](#)]
- Hoffmann, M., Kleine-Weber, H., Schroeder, S., Krüger, N., Herrler, T. and Erichsen, S. (2020). SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell*. 245-250. [[Crossref](#)]
- Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J. and Hu, Y. (2020). Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*, 395:497-506. [[Crossref](#)]
- Inès, M., Mouna, B., Marwa, E., & Dhoha, G. (2023). Biosurfactants as emerging substitutes of their

- synthetic counterpart in detergent formula: efficiency and environmental friendly. *Journal of Polymers and the Environment*, 1-13. [\[Crossref\]](#)
- Karnwal, A., Shrivastava, S., Al-Tawaha, A. R. M. S., Kumar, G., Singh, R., Kumar, A. and Malik, T. (2023). Microbial Biosurfactant as an Alternate to Chemical Surfactants for Application in Cosmetics Industries in Personal and Skin Care Products: A Critical Review. *Bio Med Research International*. [\[Crossref\]](#)
- Kişla, D., Gökmen, G. G., Evrendilek, G., Akan, T., Vlčko, T., Kulawik, P. and Özoğul, F. (2023). Recent developments in antimicrobial surface coatings: Various deposition techniques with nanosized particles, their application and environmental concerns. *Trends in Food Science and Technology*. [\[Crossref\]](#)
- Kracht, M.A., Rokos, H., Özel, M., Kowall, M., Pauli, G. and Vater, J. (1999). Anti-viral and hemolytic activities of surfactin isoforms and their methyl ester derivatives. *Journal of Antibiotics*, 52:613-619. [\[Crossref\]](#)
- Liu, W. and Li, H. (2020) COVID-19: attacks the 1-beta chain of hemoglobin and captures the porphyrin to inhibit humanheme metabolism. Preprint Revised On 2020:10.
- Mahalaxmi, I., Kaavya, J., Mohana, D.S. and Balachandar, V. (2020): COVID19 and olfactory dysfunction: a possible associative approach towards neuro degenerative diseases. *Journal of Cell Physiology*, 1-8. [\[Crossref\]](#)
- Morita, Y., Tadokoro, S., Sasai, M., Kitamoto, D. and Hirashima, N. (2011). Biosurfactant mannosyl-erythritol lipid inhibits secretion of inflammatory mediators from RBL-2H3 cells. *BBA-GenSubjects*, 1810:1302-1308. [\[Crossref\]](#)
- Mueller, C.M., Lin, Y., Viterbo, D., Pierre, J., Murray, S.A. and Shah, V. (2006). Sphorolipid treatment decreases inflammatory cytokine expression in an invitro model of experimental sepsis. A204. [\[Crossref\]](#)
- Nakanishi, M., Inoh, Y., Kitamoto, D. and Furuno, T. (2009). Nano vectors with a biosurfactant for gene transfection and drug delivery. *Journal of Drug Delivery, Science and Technology*, 5:411-420. [\[Crossref\]](#)
- Naughton, P.J., Marchant, R., Naughton, V. and Banat, I.M. (2019). Microbial biosurfactants: current trends and applications in agricultural and biomedical industries. *Journal of Applied Microbiology*, 127: 12-28. [\[Crossref\]](#)
- Nitschke, M. and Marangon, C. A. (2022). Microbial surfactants in nanotechnology: recent trends and applications. *Critical Reviews in Biotechnology*, 42(2), 294-310. [\[Crossref\]](#)
- Park, S.Y., Kim, J-H., Lee, S.J. and Kim, Y. (2013). Involvement of PKA and HO-1 signaling in anti-inflammatory effects of surfactin in BV-2 microglial cells. *Toxicology and Applied Pharmacology*, 268:68-78. [\[Crossref\]](#)
- Patel, P., Patel, R., Mukherjee, A., & Munshi, N. S. (2023). *Microbial biosurfactants for green agricultural technology*. In Sustainable Agriculture Reviews 60: Microbial Processes in Agriculture (pp. 389-413). Cham: Springer Nature Switzerland. [\[Crossref\]](#)
- Paulino, B.N., Pessoa, M.G., Mano, M.C.R., Molina, G., Neri-Numa, I.A, Pastore, G.M. (2016) Current status in biotechnological production and applications of glycolipid biosurfactants. *Applied Microbiology and Biotechnology*, 100:10265-10293. [\[Crossref\]](#)
- Perfumo, A., Rudden, M., Marchant, R. and Banat, I. (2017). *Biodiversity of biosurfactants and roles in enhancing the (bio) availability of hydrophobic substrates*. In Cellular ecophysiology of microbe, Handbook of hydrocarbon and lipid microbiology. Springer; 1-29. [\[Crossref\]](#)
- Ramasamy, S., Maheswari, P., Kavitha, P., Ravichandran, M., Sas, B. and Ramchand, C. (2009). Effect of Bacillus subtilis PB6, a natural probiotic on colon mucosal inflammation and plasma cytokines levels in inflammatory bowel disease. *Indian Journal of Biochemistry and Biophysics*, 46:79-85.
- Rodrigues, L., Banat, I.M., Teixeira, J. and Oliveira, R.. (2006). Biosurfactants: potential applications in medicine. *Journal Antimicrobial Chemotherapy*, 57: 609-618. [\[Crossref\]](#)
- Saimmai, A, Riansa-ngawong, W., Maneerat, S. and Dikit, P. (2020). Application of biosurfactants in the medical field. *World Journal of Science and Technology*, 17:154-166. [\[Crossref\]](#)
- Sajid, M.A., Singh, M.S., Cameotra, S., AlThubiani, A. (2020). Biosurfactants: potential applications as immunomodulator drugs. *Immunology Letters*, 223:71-77 [\[Crossref\]](#)

- Santos, D.K.F., Rufino, R.D., Luna, J.M., Santos, V.A. and Sarubbo, L.A. (2016). Biosurfactants: Multifunctional biomolecules of the 21st century. *International Journal of Molecular Science*, 17. [\[Crossref\]](#)
- Santos, A.P.P., Silva, M.D.S., Costa, E.V.L., Rufino, R.D., Santos, V.A., Ramos, C.S., Saruboo, L.A. and Porto, A.L.F. (2017). Production and characterization of a biosurfactant produced by *Streptomyces* sp. DPUA 1559 isolated from lichens of the Amazon region. *Brazilian Journal Medical and Biological Research*, 51:6657-6657. [\[Crossref\]](#)
- Sarangi, M. K., Padhi, S., Patel, L. D., Rath, G., Nanda, S. S., & Yi, D. K. (2022). Theranostic efficiency of biosurfactants against COVID-19 and similar viruses-A review. *Journal of Drug Delivery Science and Technology*, 103764. [\[Crossref\]](#)
- Shah, V., Doncel, G., Seyoum, T., Eaton, K., Zalenskaya, I., Hagver, R. (2005). Sphorolipids: novel glycolipid preventive agents for conception and sexual transmission. *Antimicrobial Agents and Chemotherapy*, 49: 4093-4100. [\[Crossref\]](#)
- Sil, M., Mitra, S., & Goswami, A. (2023). Probiotics and immunity: An overview. *Viral, Parasitic, Bacterial, and Fungal Infections*, 847-861. [\[Crossref\]](#)
- Singh, A. S. (2023). 15 Recent Advancements in Environmental Biotechnology: Bio surfactant is a Potent Bioremediation Method for Heavy Metals. *Applications of Environmental Biotechnology for Global Sustainability*, 112.
- Singh, P. and Cameotra, S.S. (2004). Potential applications of microbial surfactants in biomedical sciences. *Trends in Biotechnology*, 22:142-146. [\[Crossref\]](#)
- Sokhela, S., Lalla-Edward, S., Siedner, M. J., Majam, M., & Venter, W. D. F. (2023). Roadmap for achieving universal antiretroviral treatment. *Annual Review of Pharmacology and Toxicology*, 63, 99-117. [\[Crossref\]](#)
- Subramaniam, M. D., Venkatesan, D., Iyer, M., Subbarayan, S., Govindasami, V., Roy, A. & Vellingiri, B. (2020). Biosurfactants and anti-inflammatory activity: A potential new approach towards COVID-19. *Current Opinion in Environmental Science and Health*, 17, 72-81. [\[Crossref\]](#)
- Takeda, T., Sasai, M., Adachi, Y., Ohnishi, K., Fujisawa, J. and Izawa, S. (2017). Potential role of heme metabolism in the inducible expression of heme oxygenase-1. *BBA - General Subject*, 1861:1813-1824. [\[Crossref\]](#)
- Tripathy, D. B., Bhati, K., & Gupta, A. (2023). *Role of Surfactants against Covid-19: A Scientific Approach*. In *Macromolecular Symposia* (Vol. 407, No. 1, p. 2100415). [\[Crossref\]](#)
- Ukaegbu, C. I., Shah, S. R., Alara, R. O., & Thonda, O. A. (2023). *Biosurfactants as Potential Antitumor Agents*. In *Advancements in Biosurfactants Research* (pp. 439-460). Cham: Springer International Publishing. [\[Crossref\]](#)
- Vakil, H., Sethi, S., Fu, S., Stanek, A., Wallner, S., Gross, R. (2010). Sphorolipids decrease pulmonary inflammation in a mouse asthma model. *Nature*, 90. 392.
- Vanreppelen, G., Wuyts, J., Van Dijck, P., and Vandecruys, P. (2023). Sources of Antifungal Drugs. *Journal of Fungi*, 9(2), 171. [\[Crossref\]](#)
- Venugopal, A., Ganesan, H., Raja, S. S. S., Govindasamy, V., Arunachalam, M., Narayanasamy, A. & Vellingiri, B. (2020). Novel wastewater surveillance strategy for early detection of coronavirus disease 2019 hotspots. *Current Opinion in Environmental Science & Health*, 17, 8-13. [\[Crossref\]](#)
- Vollenbroich, D., Ozel, M., Vater, J., Kamp, R.M. and Pauli, G. (1997). Mechanism of inactivation of enveloped viruses by the biosurfactant surfactin from *Bacillus subtilis*. *Biologicals*, 25: 289-297. [\[Crossref\]](#)
- Wang, W., He, J. and Wu, S. (2020). The definition and risks of cytokine release syndrome-like in 11 COVID-19-infected pneumonia critically ill patients: disease characteristics and retrospective analysis. *Medrxiv*. ppzbxmed-10.1101.2020.02.26.20026989
- Yang, M. (2020). Cell pyroptosis, a potential pathogenic mechanism of 2019-nCoV infection. Available at: SSRN3527420 2020. This article was published, and it explains the SARS-CoV-2 outbreak, treatment and other related approaches towards COVID-19. [\[Crossref\]](#)
- Zhang, Y., Liu, C., Dong, B., Ma, X., Hou, L. and Cao, X. (2015). Anti-inflammatory activity and mechanism of surfactin in lipopolysaccharide-activated macrophages. *Inflammation*, 38:756-764. [\[Crossref\]](#)