





## ORIGINAL RESEARCH ARTICLE

## Phytochemical screening and antioxidant activities of the bark of *Lawsonia inermis* (Henna) grown in Dekina, Kogi, Nigeria

Abuh Omachoko Leonard<sup>1\*</sup> , Isah Aminu Agih<sup>2</sup> , Azeez Ayinla Akeem<sup>3</sup> , and Kwasi Ndanusa Benjamin.<sup>4</sup> 

<sup>1</sup>Department of Integrated Science, Kogi State College of Education, Ankpa, Nigeria

<sup>2,3,4</sup>Department of Chemistry, Kogi State College of Education Ankpa, Nigeria

### ABSTRACT

Severe health conditions can result from lack of knowledge on the bioactive constituents of medicinal plants, as it the main stay of treatment and prevention of diseases in our locality. Hence, the need to examine the therapeutic active ingredients responsible for plants medicinal ability for better health management. This study analyses the phytochemical constituents and anti – oxidant activity of the bark of *lawsonia inermis*. *Lawsonia inermis* bark was collected, cut into pieces and dried at room temperature. The dried sample was pulverized into fine particle. 200g of the pulverized sample was weighed into a 750mL container and was added 300mL absolute ethanol, then allowed to stand for extraction. The solute obtained by evaporating the ethanol was screened for the presence of phyto – components and the result shows alkaloids, glycosides, saponins, phenols, flavonoids, proteins and amino acid as bioactive substances. Also, *lawsonia inermis* bark displayed an efficient free radical scavenging potential as the anti – oxidant ability showed the percentage KMnO<sub>4</sub> radical scavenging activities of 41.4% which shows close competence with the standard ascorbic acid; 47.9% at EC<sub>50</sub>; This indicate *lawsonia inermis* bark to exhibit a good potential as antioxidant and could scavenge free radical from the body. The presence of the bioactive substances signifies its usefulness in therapeutic health as agent that could inhibit diseases causing several ailments and as well be a good source for health management control.

### ARTICLE HISTORY

Received August 10, 2022

Accepted September 27, 2022

Published on September 30, 2022

### KEYWORDS

Phytochemical, screening, *lawsonia inermis*, antioxidant

© The authors. This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>)

### INTRODUCTION

Management of plants in disease treatment and prevention have been in use over a long period of time, i.e. since ages (Mohan, *et al.*, 2021), with communities been ignorant of the active ingredient responsible for the potency. Plants are used in the treatment of diseases like hypertension, pile, typhoid fever, asthma, rheumatism etc. (Mohan, *et al.*, 2021; Farzama, *et al.*, 2021 and Nguyen, *et al.*, 2021). Medicinal plants are known to possess the richest bio resources such as anti-neoplastic, antimicrobial, antioxidant, anti-inflammatory etc. (Junaid, *et al.*, 2020; Farzama, *et al.*, 2021; Mohan, *et al.*, 2021; Letiele, *et al.*, 2021; Isah, *et al.*, 2020; Nigam & Arnold, 2020). Their medicinal potential is a result of natural secondary metabolites which include flavonoids, saponins, tannins, alkaloids, phenols etc. that fight against bacterial infections and related diseases (Farzama, *et al.*, 2021). These secondary metabolites are known as phytochemicals and they work in conjunction with nutrients and dietary fiber to protect the body from diseases and infections; they also contribute to

regeneration of cells in the body (Isah, *et al.* 2020). Plants inherent ability to biosynthesize a diverse range of non-enzymatic antioxidants capable of mitigating oxidative damage is attributed to low progression of several diseases caused by free radicals (Deepak, *et al.*, 2015), they also have evolved a variety of active substances that scavenge free radicals and thus protect cells, delay aging and protect diseases associated with aging (Hassan, *et al.* 2017), these substances can significantly reduce or inhibit the rate of oxidation and participate in physiological processes and also terminate chain reaction caused by free radicals that are linked to diseases like cancer, arthritis, liver damage, and diabetes (Olamide, *et al.*, 2017; Shohber, *et al.*, 2015). The role of plants as antioxidant have shown its ability to remove free radicals from nerves and thus effectively protect against most nervous system diseases. Bioactive substances such as polyphenols, vitamins, alkaloids, polysaccharides, and active peptides in plants aids in the preservation of neuron structure and the maintenance of a healthy state

**Correspondence:** Abuh Omachoko Leonard. Department of Integrated Science, Kogi State College of Education, Ankpa, Nigeria. ✉ [leonardabuh@gmail.com](mailto:leonardabuh@gmail.com), Phone: 08036002152

**How to cite:** Abuh Omachoko Leonard, Isah Aminu Agih, Azeez Ayinla Akeem, and Kwasi Ndanusa Benjamin. (2022). Phytochemical screening and antioxidant activities of the bark of *Lawsonia inermis* (Henna) grown in Dekina, Kogi, Nigeria.. *UMYU Scientifica*, 1(2), 274 – 279. <https://doi.org/10.56919/usci.1122.035>

have been the major sources of antioxidants used in reducing oxidative stress (Muhamat, et al. 2021). These potentials are also attributed to *Lawsonia inermis*.

*Lawsonia inermis* is commonly known as henna, it belongs to the kingdom plantae, phylum; spermatophyte, sub phylum; angiosperm, class; dicotyledon, order; myrtales and family; lythraceae (Rojas – Sandovaj (2017). It is a plant that have been part of human existence for curbing many diseases. Researches conducted by scholars indicted the significant medicinal importance of *Lawsonia inermis* in the treatment and management of diseases such as leprosy, fever, leucorrhoea, rheumatoid, arthritis, ulcer, cardiac disease and headache (Arirudran, et al., 2019; Chetty, 2008; Reddy, 1988). The medicinal value of *Lawsonia inermis* also found significance in the treatment of wounds, blood sugar, microbial infections, inflammations and proliferation of cancer cells (Debrayiya, et al., 2020). The healing properties like analgesic, hypoglycemic, hypotoprotive, immuno stimulant, anti-inflammatory, antimicrobial, antibacterial, antifungal, antiviral anticancer, anti-parasitic, anti- dermatophytic properties etc. were attributed to the presence of phytochemicals components like carbohydrate, proteins, flavonoids, tannins and phenolic compounds (Gagandeep, et al., 2010). Phytochemicals substances; flavonoids, steroids, tannins, protein, cardioglycosides, alkaloids and carbohydrate were reported on the flowers of *lawsonia inermis* to be responsible for its therapeutic potentials (Syede, 2018). The radical scavenging ability denotes the inhibition potential of the plant bark with strong antioxidant capabilities and medicinal role in combating diseases causing different ailment (Dharmesh, et al., 2020). Traditional medicine is in constant use as a natural remedy in our locality for the treatment of several ailments, with the community having little or no knowledge of the active ingredients responsible for its medicinal potency, hence, the need to elucidate the phytochemical components of this plant part. Despite the important role of *Lawsonia inermis* as a source of natural antioxidant in disease treatment and health management in our society, hazard attributed to drug toxicity that could lead to severe health conditions and possible death could arise as a result of overdose and basically lack of understanding of the phyto – constituents. This study is aimed at investigating the phytochemical substances and antioxidant activity of the bark of *Lawsonia inermis*.

## MATERIALS AND METHODS

The materials used include pulverized *lawsonia inermis* bark which was extracted using absolute ethanol (99%), sodium hydroxide, ammonia solution, hydrochloric acid, ferric chloride solution, concentrated sulphuric acid, nitric acid, Hager reagent, Gelatin solution, sodium nitroprusside, pyridine, Benedict solution were standard analytical reagents used for the phytochemicals analysis.  $\text{KMnO}_4$ , Ascorbic acid, Hydrogen peroxide and double beam UV / Vis spectrophotometer 2700

## Collection and sample Treatment

Bark of *Lawsonia inermis* was collected in Dekina, Kogi State and was identified and authenticated by a botanist, Professor Simon Sugei Usman and the sample deposited at our institution department of Biology. It was cut into pieces and air dried at room temperature. The plant part was then finely ground. 200g of the pulverized sample was weighed into a 750mL plastic container, followed by 300mL of absolute ethanol and corking. For proper extraction, the mixture was left for a week. The extract was filtered, and the dried solute was sacrificially obtained (Leonard, et al., 2022).

## Phytochemical screening

Phytochemical screening method employed are as described by Prashant et al. (2011); Junaid & Pati (2020).

### Test for alkaloids

*Hager's test*: the extract (2g) was weighed and dissolved in 5mL diluted HCl and filtered. 1mL of Hager's reagent was mixed with 2mL of the filtrate. The presence of alkaloids will be confirmed by a creamy white precipitate.

### Test for saponins

*Foam test*: sample of the extract, (1g) was dissolved in distilled water and vigorously shaken before being allowed to stand for 10 minutes. Saponins will be detected by a stable persistent foam.

### Test for phytosterols

*Salkowski's test*: the extract was weighed (2g) and treated with chloroform and then filtered; the filtrate was then treated with a few drops of concentrated  $\text{H}_2\text{SO}_4$ ; the mixture was shaken and allowed to stand until a golden yellow color appeared.

### Test for flavonoids

*Alkaline reagent test*: few drops of sodium hydroxide solution was added to 2g of the extract, formation of an intense yellow color that will turn colorless upon addition of dilute HCl acid solution will confirm the presence of flavonoids.

### Test for carbohydrate

*Benedict's test*: the plant extract, (2g) was weighed and 5mL distilled water was added and dissolved, then filtered. A 0.2 mL sample of the filtrate was treated with Benedict reagent and gently heated. The presence of carbohydrate will be confirmed by the formation of orange-red precipitate.

### Test for tannins

*Gelatin test*: the extract (0.2g) was weighed and dissolved in 5mL of distilled water, followed by 1% gelatin solution and 10% sodium chloride solution. The presence of tannins will be indicated by a white precipitate.

*Legal's test*: the plant extract, (0.5g) was weighed and dissolved in pyridine, sodium nitroprusside, and 10%

sodium hydroxide. When glycosides are present, the solution will turn pink.

**Test for glycosides**

*Legal's test:* the plant extract, (0.5g) was weighed and dissolved in pyridine, sodium nitroprusside, and 10% sodium hydroxide. When glycosides are present, the solution will turn pink.

**Test for phenols**

*Ferric chloride test:* the plant extract, (0.2g) was treated with 3-4 drops of ferric chloride solution. The presence of phenols will be indicated by the formation of a bluish-black color.

**Test for protein and amino acids:**

*Xanthoproteic test:* the plant extract, (1g) was treated with few drops of concentrated nitric acid. Formation of yellow color will indicate the presence of protein.

**Antioxidant Activities**

Radical scavenging method using acidified potassium permanganate (KMnO<sub>4</sub>) was used. Ascorbic acid at concentration of 2.0mg/mL – 0.25mg/mL (I e 2.0, 1.5, 1.0, 0.5 & 0.25mL) was used as standard antioxidant. 0.5mL of acidified potassium permanganate solution and 0.5mL of hydrogen peroxide was added as pro oxidant. Absorbance of both the KMnO<sub>4</sub>/ascorbic acid and KMnO<sub>4</sub>/sample extract were taken spectrophotometrically at a wavelength of 520nm after 30 minutes' incubation period. The graph of absorbance against concentration was then plotted. The % KMnO<sub>4</sub> scavenging effect was determined using

$$\% \text{KMnO}_4 \text{ scavenging effect} = \frac{Ac - At}{Ac} * 100/1$$

Where:

Ac = Absorbance of control

At = Absorbance of the tested sample (Isaac *et al.*, 2011)

**Preparation of standard solutions**

*Ascorbic acid:* (0.011 mol/ 100mL),

0.2g of ascorbic acid was weighed and dissolved in 100mL of distilled water.

*Potassium permanganate:* (0.0006 mol/100mL)

0.01g of potassium permanganate was weighed and dissolved in 100mL of distilled water to make the stock, from the stock 25mL of KMnO<sub>4</sub> solution was acidified by adding 10mL of 2M H<sub>2</sub>SO<sub>4</sub> solution.

**RESULTS**

**Phytochemical screening of *lawsonia inermis* bark**

The phytochemical substances identified are alkaloids, glycosides, saponins, phenols, flavonoids, protein and amino acid, while Carbohydrates, phytosterols and tannins were absent as shown in Table 1.


**Antioxidant activity using %KMnO<sub>4</sub> scavenging effect**

The % KMnO<sub>4</sub> scavenging effect for the sample is 41.4% and that of the standard is 47.9% at EC<sub>50</sub>. Fig.1 below shows the graphical competence of the bark of *lawsonia inermis* and indicate its antioxidant properties.

**Table 1: Phytochemical screening Result**

Parameters	Test performed	Observation	Results
Alkaloids	Hager test	Creamy white precipitate formed	+
Carbohydrates	Benedicts test	Orange – red precipitate formed	-
Glycosides	Legal test	Pink solution formed	+
Saponins	Foam test	Persistent foam formed	+
Phytosterols	Salkowskis test	Golden yellow solution formed	-
Phenols	Ferric chloride test	Bluish – black colour formed	+
Tannins	Gelatin test	White precipitate formed	-
Flavonoids	Alkaline test	Intense yellow color turns colorless in dil HCl	+
Protein and amino acid	Xanthoproteic test	Yellow color formed	+

**Key:** (+) = positive; (-) = absent



**Figure 1:** Result of the antioxidants activities of *Lawsonia Inermis* bark

## DISCUSSION

Phytochemical screening result of this study show the therapeutic potential of the bark of *Lawsonia inermis* plant as a result of the presence of bioactive substances in it. Previous work conducted on this plant also present bioactive ingredients, although with some variation from the current study. A work conducted at Lahore, Pakistan by Iram *et al.* (2013) indicted the presences of cardioglycosides, terpenoids, carbonhydrate, phenols, quinones and tannins with the absent of protein on the methanol, acetone and aqueous extract of the leaves. Nirmala *et al.* (2016) reported the methanol extract to contain carbohydrate, protein, flavonoids, tannins, phenolic compounds, alkaloids, terpenoids, glycoside, saponins, quinones, protein and amino acid in a study they conducted at Telegona, India. Gabriel *et al.* (2017) reported a quantitative presence of total alkaloids, total flavonoids, total phenol, total saponin and total tannin with glycosides and protein been absent in a research they carried out on the ethanolic extract of the stem bark of *Lawsonia inermis* in Ogun, western Nigeria. In another work by Dharmesh *et al.* (2020) in india, flavonoids, polyphenols, tannins, and terpenoids were reported to be present in the methanol extract of the bark of *lawsonia inermis* plant with alkaloids, saponins, proteins and glycoside to be absent.

The variations in the presence of the bioactive constituents as seen in the results could be as a factor of the extracting solvents used, as different extracting solvent possesses different polarities. Also, phyto - substituents are related to their geographical origins; the geographical location or origin of plants as reported by Mohammed and Aisha (2022) could contribute to the variation in the phyto – component present in the plant. The antioxidant predicts the plant bark to possess a great potential to scavenge free radical, this finding agrees with Debapriya *et al.* (2020) in a work they uphold the radical scavenging potentials of

*Lawsonia inermis* in treating human disease like cancer. In a similar vein, Dharmesh *et al.*, (2020) affirm the anti – oxidative activity of the plant in their study where *Lawsonia inermis* bark indicted inhibitory capacity in DPPH, Superoxide and Ferric chloride methods of free radical scavenging activity.

## CONCLUSION

Plants have been in use as medicine over decades in the treatment of different kind of illnesses; despite the advancement in modern medicine, traditional medicine which greatly utilizes plants is still gaining significance; this is so because of its preventive and curative potential which is as a result of the presence of bioactive substances. This study has shown the presence of alkaloids, glycosides, saponins, phenols, flavonoids, protein and amino acid as bioactive substances. They phyto – chemical constituents and the efficient free radical scavenging ability displayed by the bark of *Lawsonia inermis* indicates that it could inhibit oxidation process and possibly prevent oxidative damage on cells and tissues. This study affirms the presence of therapeutic substances responsible for its efficacy when use as traditional medicine. The plant possesses medicinal properties and if properly exploited, it can potentially be used to treat ailment caused by free radicals and also it could significantly serve as substitute agent in therapeutics. However, standardization and isolation into pure form and the investigation of the toxicity level is recommended for better and efficient health management control.

## ACKNOWLEDGEMENT

We acknowledge the laboratory technologist Mr David Agih of the department of Chemistry and Mallam Ogagi Adams of the department of Biology, Kogi State College of Education Ankpa in providing all the necessary materials and support for the success of this work.

**REFERENCES**

- Arirudran, V., Krishnan, U. S. and Mahadeva, R. (2019). Preliminary phytochemical analysis and isolation and identification of active principle component of petroleum extract from leaves of lawsonia inermis L. *Research journal of pharmacy and technology*, 12(8), 3629 - 3636. [\[Crossref\]](#)
- Chetty, K. M. (2008). Flowering of Chittor. Edn 1. Andhra Pradesh. Pp132
- Debabriya, D., Dipu, S., Rajat, B., Suchita, S., Bidisha, M., Sayak, G. and Debleenn, R. (2020). Insight into the phytochemical potential of lawsonia inermis for future small molecule based therapeutic application. *International research journal*. 11(1), 1 - 7. [\[Crossref\]](#)
- Deepak, M. K., Srendra, S. K., Mahabaleshwar, V. H. and Hanhong, B. (2015). Significance of antioxidant potential of plants and its relevance to therapeutic application. *Int. J. Biol Sci*, 11(8), 982 - 991. [\[Crossref\]](#)
- Del Rio, L. A. (2015). ROS and RNS in plant physiology: an overview. *J. Exp. Bot.* 66, 22827 – 2837. [\[Crossref\]](#)
- Dharmesh, G., Darpini, S. P., Kishor, K. D., Santosh, S and Timir, B. P. (2020). Phytochemical and antimicrobial activity methanolic extract of lawsonis inermis L bark. *Current issues in pharmacy and medical sciences*. 33(3). 132 - 138. [\[Crossref\]](#)
- Farzana, K., Mohammed - Awais, F., Hafiz - Muhammad, U. F., Abdul - Rahim, S., Atif - Ali, K. K., Chul - Woong, K., Muhamma, H. M., Gaber- El- saber, B., Kyung, H.C. and Abdul, S.M. (2021). Phytochemical Investigation, Antimicrobial, Antioxidant and Anticancer Activities of *Acer Cappad Ocicum* *Gid*, doi.org/10.3390.life.11070656.
- Gabriel, A. A., Olukayode, Y. O. and Taiwo, O. E. (2017). Quantitations of phytochemicals and biocide actions of lawsonia inermis Linn extract against wood termites and fungi. *International Biodeterioration and Biodegradation*. 116. 155 - 162. [\[Crossref\]](#)
- Gagandeep, C., Sandeep, G. and Priyanka, P. (2010). Lawwsonia inermis Linnaeus: a phytopharmacological Review. *International journal of pharmaceutical sciences and drug Research*. 2 (2). 91 - 98
- Hassan, W., Noreen, H., Rehman, S., Gul, S., Kamal, M. A., Kamdem, J. P. (2017). Oxidative stress and antioxidant potential of one hundred medicinal plants. *Curr.Top.Med.chem*, 17, 1336 - 1370. [\[Crossref\]](#)
- Iram, G., Maria, S., Muhammed, S. A. and Muhammed, A. A. (2013). Phytochemical, toxicological and antimicrobial evaluation of lawsonia inermis extract against clinical isolation of pathogenic bacteria. *Ann. Clin microbial antimicrobs*. 12. 36. [\[Crossref\]](#)
- Isaac, K. A., Emmanuel, O., Abraham, Y. M., Francis, M. S., Francis, A. and Linda, M. S. (2016). Development and validation of Radical Scavenging antioxidant assay using potassium permanganate. *J. Scientific & Innov. Res*, 5(2), 36-42. [\[Crossref\]](#)
- Isah, A. A., Abuh, L. O., Omachi, A. A, Usman, A. and Azeez, A. A. (2020). proximate analysis, Phytochemical screening, Antioxidant activities and Potassium/Sodium determination of yellow Ipomoea batata grown in Ankpa, Kogi state Nigeria. *FUW trends in science & technology journal*, 5(1), 226 - 230.
- Jiraporn, L., Churdsak, J., Kanokporn, S. and Paiwan, S. (2021). Phytochemical screening, Antioxidant and sperm Viability of *Nelumbo nucifera* petal extract. *MDPI*, 10(7), 1375. [\[Crossref\]](#)
- Junaid, R. S. and Patil, M. K. (2020). Qualitative test for preliminary phytochemical screening; An Overview. *International Journal of Chemical studies*, 8(2), 603 - 608. [\[Crossref\]](#)
- Lancho, M. M., Getachew, M. D., Dagmawat, A. B., Edilu, J. S, Askale, G., Wakuma, M. B., Morka, D. B., Petros, A., Abraham, M., Miressa, T., Kebede, A., and Dejene, B. (2021). Phytochemical screening and in - Vitro evaluation of Antibacterial activities of *Echinops amplexicaulis*, *Ruta Chalepenasis* and *salix subserrata* against selected

- pathogenic bacterial strains in West Shewa zone, Ethiopia. *Journal of Experimental pharmacol* 13, 511 - 520. [\[Crossref\]](#)
- Leonard, O. A. Salifu, V. and Samuel, A. S. (2022). Dyestuff utilization of the bark of parkia biglobosa grown in Ankpa, Kogi, Nigeria. *FUW Trend in science and technology journal*.7(2). 862 - 866.
- Letiele, B. D., Amanda, L. G., Thainara, D. A., Thais, F. K., Jefferson, E., Rafael, R. M., Vanusa, M. and Elton, I. G. D. (2020). phytochemical analysis, Antioxidant Activity, Antimicrobial Activity and Cytotoxicity of Chaptalia Nutans leaves. *Hindawii Advance in Pharmacological and pharmaceutical Science*, [\[Crossref\]](#)
- Mohammed, S. A. and Aisha, A. Y. (2022). Comparative phytochemical screening and acute toxicity study of two varieties of ginger, zingiber officinale. *UMYU Scientifica*. 1(1). 6 - 11. [\[Crossref\]](#)
- Mohan, E., Suniya, S., Shanmugan, S. and Rajendran, K. (2021). Qualitative Phytochemical screening of selected medicinal plants. *Journal of Drug Delivery and Therapeutic*, 11(2), 141- 144. [\[Crossref\]](#)
- Muhamat, S. A., Lahat, N., Cyrille, A. N., Mady, C. and Mar, D. C. (2021). Phytochemical screening, Total Phenolic and Antioxidant activities of numerous extracts from leaves and bark of bauhinia rufescens lam. *Food and Nutrition Science*,12(6), 544 - 551. [\[Crossref\]](#)
- Nguyen, T. M., Dang, X. C., Pham, V. T., Trying, N. M., Thuc- Huy, D., Tran, T. L. M. and Vo, T. T. (2021). Phytochemical screening and Evaluation of Antioxidant properties and Antimicrobial activities against xanthomonas axonopodis of Euphorbia tricucalli extract in Binh Thuan Province, Vietnam. *MDPI Molecules*, doi.org/10:3390/Molecules. 26040941.
- Nigam, R. and Arnold, R. (2020). Qualitative and Quantitative Phytochemical screening and Clinical finger print Analysis of Herbal plant phyllanthus niruri using HPTLC. *Journal of Scientific Research*, 13(2), 623 - 633. [\[Crossref\]](#)
- Nirmala, B.R., Sitakumari, O. and Rajesh, G. G. (2016). Phytochemical analysis and antimicrobial activity of Lawsonis inermis (Henna). *Journal of plant science and research*. 3(2). 158.
- Olamide, E. A., Funsho, O. O., Tan, N., and Zeng, G. (2017). In Vitro antioxidant activity, total phenolic and flavonoid contents of ethanol extract of stem and leaf of Grewia carpinifolia. *Beni - Suef university journal of Basic and Applied Sciences*, 6(1), [\[Crossref\]](#)
- Prashant, T., Bimlesh, K., Mandeep, K., Gurpreet, K. and Harleen, K. (2011). Phytochemical screening and Extraction: A review. *International Pharmaceutical Scientia*, 1(1), 98 - 106.
- Reddy, K. R. (1988). Folk medicine from Chittor district Andhra Pradesh India used in the treating of jaundice. *International journal of crude drug research*. 26. 137 - 140. [\[Crossref\]](#)
- Rojas-Sandovaj, J. (2017). *Lawsonia inermis* (Egyptian privet). *Invasive species compendium*. wallingford. Uk. CABI. [\[Crossref\]](#)
- Shobha, P. R., Shanthi, V., and Rashed, M. A. (2015). Evaluation of Antioxidant activities of some medicinal plants using oxidizing agents in acidic medium by colorimetry. *Int. J. Innov. Sci. Engr & Techn*. 2(5). Retrieved from [www.ijiset.com](http://www.ijiset.com) on 12/04/2022
- Syede, N. F. (2018). Pharmacognostic assessment of Lawsonia inermis flowers. *Journal of pharmacognosy and phytochemistry*. 7(6). 2365 - 2369